

Terreni di coltura

La Biotoxik è lieta di presentare una vasta gamma di prodotti di marchio Biotec utili nel QC laboratorio delle industrie farmaceutiche, cosmetiche e alimentari.

Confezionamento con terzo involucro di cartone: prezzo: Euro 1,00 da aggiungere al prezzo

| Culture media | | | | | | |
|--|--|-------|---------------------------------|------------|------|-------|
| Product Name | Use | Code | Packaging | Shelf Life | °C | Price |
| A1 MEDIUM | For the determination of Coliforms in water and food | 20848 | 10 glass tubes | 180 days | 4-8 | |
| A1 MEDIUM | For the confirmation of Pseudomonas aeruginosa | 6184 | Dehydrated 500 gr | 3 years | 8-25 | |
| ACETAMIDE AGAR | For the confirmation of Pseudomonas aeruginosa | 6267 | Dehydrated 500 gr | 3 years | 8-25 | |
| ACETAMIDE BROTH ISO 16266 | For the confirmation of Pseudomonas aeruginosa | 20041 | 10 glass tubes 5 ml | 90 days | 4-8 | |
| ACETAMIDE BROTH BASE ISO 16266 | For the confirmation of Pseudomonas aeruginosa | 6193 | Dehydrated 500 gr | 3 years | 8-25 | |
| ACETAMIDE BROTH BASE ISO 16266 – REQUIRED SUPPLEMENT | For the confirmation of Pseudomonas aeruginosa | 6532 | 5 eppendorf x 1 lt / B Solution | 90 days | 4-8 | |
| ACETAMIDE BROTH | For the confirmation of Pseudomonas aeruginosa | 6185 | Dehydrated 500 gr | 3 years | 8-25 | |
| ACETOBACTER | For the determination of Acetobacter spp. | 20395 | 20 plates 90 mm | 180 days | 4-8 | |
| ACETOBACTER | For the determination of Acetobacter spp. | 20522 | 40 plates 60 mm | 180 days | 4-8 | |
| ACETOBACTER | For the determination of Acetobacter spp. | 6803 | Dehydrated 500 gr | 3 years | 8-25 | |
| ACID BUFFER ISO 11731 | For the washing of the filter membrane in the detection of Legionella spp. | 20621 | 4 bottles 100 ml | 270 days | 8-25 | |
| ACID BUFFER ISO 11731 | For the washing of the filter membrane in the detection of Legionella spp. | 20709 | 10 glass tubes 9 ml | 270 days | 8-25 | |
| ACID BUFFER ISO 11731 | For the washing of the filter membrane in the detection of Legionella spp. | 20728 | 4 bottles 30 ml | 270 days | 8-25 | |
| ACID BUFFER ISO 11731 | For the washing of the filter membrane in the detection of Legionella spp. | 20966 | 4 bottles 30 ml 10X | 270 days | 8-25 | |
| ACID BUFFER ISO 11731 | For the washing of the filter membrane in the detection of Legionella spp. | 20965 | 4 bottles 100 ml 10X | 270 days | 8-25 | |
| AEROMONAS HYDROPHILA (RYAN) UNICHIM 1039:2002 | For the isolation of Aeromonas spp. | 20635 | 20 plates 90 mm | 180 days | 4-8 | |
| AEROMONAS AGAR BASE (RYAN) UNICHIM 1039:2002 | For the isolation of Aeromonas spp. | 6269 | Dehydrated 500 gr | 3 years | 8-25 | |
| AEROMONAS AGAR BASE (RYAN) UNICHIM 1039:2002 – REQUIRED SUPPLEMENT | | 6331 | 10 vials x 500 ml / AMPICILLIN | 3 years | 4-8 | |
| AGAR, EUROPEAN BACTERIOLOGICAL | For use in microbiology | 6401 | Dehydrated 500 gr | 3 years | 8-25 | |
| AGAR, INDUSTRIAL | For the use in the food industry | 6402 | Dehydrated 500 gr | 3 years | 8-25 | |
| ARGININA DECARBOSSILASI BROTH | For the test of the decarboxylation of the amino acids | 1875 | 20 polystyrene tubes | 240 | 4-8 | |
| AZIDE DEXTROSE BROTH (WHO-APHA) | For the selective enrichment of Enterococcus spp. in water and food | 1156 | 10 glass tubes | 270 days | 8-25 | |
| AZIDE DEXTROSE BROTH (WHO-APHA) | For the selective enrichment of Enterococcus spp. in water and food | 20667 | 10 glass tubes 9 ml | 270 days | 8-25 | |

| | | | | | | |
|---|---|-------|--|----------|------|--|
| AZIDE DEXTROSE BROTH (WHO-APHA) | For the selective enrichment of Enterococcus spp. in water and food | 6621 | Dehydrated 500 gr | 3 years | 8-25 | |
| AZIDE DEXTROSE BROTH 2X (WHO-APHA) | For the selective enrichment of Enterococcus spp. in water and food | 21198 | 10 glass tubes | 270 days | 8-25 | |
| BACILLUS CEREUS SELECTIVE (PEMBA) ISO 21871 ISTISAN 96/35 | Isolation of B. Cereus from foods and milk | 1087 | 20 plates 90 mm | 180 days | 4-8 | |
| BACILLUS CEREUS SELECTIVE (PEMBA) ISO 21871 ISTISAN 96/35 | Isolation of B. Cereus from foods and milk | 4087 | 40 contact | 180 days | 4-8 | |
| BACILLUS CEREUS SELECTIVE (PEMBA) ISO 21871 ISTISAN 96/35 | Isolation of B. Cereus from foods and milk | 2287 | 40 plates 60 mm | 180 days | 4-8 | |
| BACILLUS CEREUS (PEMBA) AGAR BASE ISO 21871 ISTISAN 96/35 | Isolation of B. Cereus from foods and milk | 6718 | Dehydrated 500 gr | 3 years | 8-25 | |
| BACILLUS CEREUS (PEMBA) AGAR BASE ISO 21871 ISTISAN 96/35 – REQUIRED SUPPLEMENT | | 6304 | 1 x 100 ml / EGG YOLK EMULSION | 180 days | 4-8 | |
| BACILLUS CEREUS (PEMBA) AGAR BASE ISO 21871 ISTISAN 96/35 – REQUIRED SUPPLEMENT | | 6315 | 10 vials x 500 ml / BACILLUS CEREUS SUPPLEMENT | 3 years | 4-8 | |
| BACILLUS CEREUS SELECTIVE (MYP) ISO 7932 | Isolation of B. Cereus from foods and milk | 20557 | 20 plates 90 mm | 180 days | 4-8 | |
| BACILLUS CEREUS SELECTIVE (MYP) ISO 7932 | Isolation of B. Cereus from foods and milk | 20605 | 40 contact | 180 days | 4-8 | |
| BACILLUS CEREUS (MYP) AGAR BASE ISO 7932 | Isolation of B. Cereus from foods and milk | 6187 | Dehydrated 500 gr | 3 years | 8-25 | |
| BACILLUS CEREUS (MYP) AGAR BASE ISO 7932 – REQUIRED SUPPLEMENT | | 6304 | 1 x 100 ml / EGG YOLK EMULSION | 180 days | 4-8 | |
| BACILLUS CEREUS (MYP) AGAR BASE ISO 7932 – REQUIRED SUPPLEMENT | | 6315 | 10 vials x 500 ml / BACILLUS CEREUS SUPPLEMENT | 3 years | 4-8 | |
| BAIRD PARKER | Isolation of S. aureus | 1074 | 20 plates 90 mm | 180 days | 4-8 | |
| BAIRD PARKER | Isolation of S. aureus | 20172 | 20 plates 90 mm 30 ml | 180 days | 4-8 | |
| BAIRD PARKER | Isolation of S. aureus | 2274 | 40 plates 60 mm | 180 days | 4-8 | |
| BAIRD PARKER | Isolation of S. aureus | 4074 | 40 contact | 180 days | 4-8 | |
| BAIRD PARKER | Isolation of S. aureus | 2574 | 40 plates s/m | 180 days | 4-8 | |
| BAIRD PARKER AGAR BASE (pH 6,8) | Isolation of S. aureus | 6003 | Dehydrated 500 gr | 3 years | 8-25 | |
| BAIRD PARKER AGAR BASE (pH 6,8) | Isolation of S. aureus | 20859 | 4 bottles 90 ml | 180 days | 8-25 | |
| BAIRD PARKER AGAR BASE (pH 6,8) – REQUIRED SUPPLEMENT | | 6310 | 1 x 100 ml / EGG YOLK EMULSION + POTASSIUM TELLURITE | 180 days | 4-8 | |
| BAIRD PARKER AGAR BASE | Isolation of coagulase positive Staphylococcus in the food | 6188 | Dehydrated 500 gr | 3 years | 8-25 | |
| BAIRD PARKER AGAR BASE | Isolation of coagulase positive Staphylococcus in the food | 20676 | 4 bottles 90 ml | 180 days | 4-8 | |
| BAIRD PARKER AGAR BASE – REQUIRED SUPPLEMENT (ISO 6888-1) | | 6310 | 1 x 100 ml / EGG YOLK EMULSION + POTASSIUM TELLURITE | 180 days | 4-8 | |
| BAIRD PARKER AGAR BASE – REQUIRED SUPPLEMENT (ISO 6888-2) | | 6341 | 10 vials x 100 ml / RPF SUPPLEMENT | 2 anni | 4-8 | |
| BAIRD PARKER ISO 6888-1 | Isolation of coagulase positive Staphylococcus in the food | 20995 | 20 plates 90 mm | 180 days | 4-8 | |
| BAIRD PARKER ISO 6888-1 | Isolation of coagulase positive Staphylococcus in the food | 20996 | 40 plates 60 mm | 180 days | 4-8 | |
| BAIRD PARKER ISO 6888-1 | Isolation of coagulase positive Staphylococcus in the food | 20997 | 40 contact 55 mm | 180 days | 4-8 | |

| | | | | | | |
|---|--|-------|---|----------|------|--|
| BAIRD PARKER RPF ISO 6888-2 | Isolation of coagulase positive <i>Staphylococcus</i> in the food | 20202 | 20 plates 90 mm | 180 days | 4-8 | |
| BAIRD PARKER RPF ISO 6888-2 | Isolation of coagulase positive <i>Staphylococcus</i> in the food | 21181 | 40 plates 60 mm | 180 days | 4-8 | |
| BAIRD PARKER RPF ISO 6888-2 | Isolation of coagulase positive <i>Staphylococcus</i> in the food | 20825 | 40 contact 55 mm | 180 days | 4-8 | |
| BCP GLUCOSE AGAR ISO 21528 e ISO 11059 | For the determination of Enterobacteria - For the confirmation of <i>Pseudomonas</i> spp | 20477 | 10 glass tubes 15 ml | 180 days | 8-25 | |
| BCP GLUCOSE AGAR ISO 21528 e ISO 11059 | For the determination of Enterobacteria - For the confirmation of <i>Pseudomonas</i> spp | 20445 | 4 bottles 100 ml | 180 days | 8-25 | |
| BCP GLUCOSE AGAR ISO 21528 e ISO 11059 | For the determination of Enterobacteria - For the confirmation of <i>Pseudomonas</i> spp | 20494 | 10 glass tubes slant 7 ml | 180 days | 8-25 | |
| BCP GLUCOSE AGAR ISO 21528 e ISO 11059 | For the determination of Enterobacteria - For the confirmation of <i>Pseudomonas</i> spp | 20744 | 20 plates 90 mm | 180 days | 8-25 | |
| BCP GLUCOSE AGAR ISO 21528 e ISO 11059 | For the determination of Enterobacteria - For the confirmation of <i>Pseudomonas</i> spp | 20371 | 40 plates 60 mm | 180 days | 8-25 | |
| BCP GLUCOSE AGAR ISO 21528 e ISO 11059 | For the determination of Enterobacteria - For the confirmation of <i>Pseudomonas</i> spp | 6190 | Dehydrated 500 gr | 3 years | 8-25 | |
| BEEF EXTRACT | For use in microbiology | 6404 | Dehydrated 500 gr | 3 years | 8-25 | |
| BILE ESCULIN AZIDE ISO 7899 APAT IRSA 29:2003 | Isolation of <i>Enterococcus</i> spp. | 1032 | 20 plates 90 mm | 270 days | 8-25 | |
| BILE ESCULIN AZIDE ISO 7899 APAT IRSA 29:2003 | Isolation of <i>Enterococcus</i> spp. | 2232 | 40 plates 60 mm | 270 days | 8-25 | |
| BILE ESCULIN AZIDE ISO 7899 APAT IRSA 29:2003 | Isolation of <i>Enterococcus</i> spp. | 1132 | 10 glass tubes slant | 270 days | 8-25 | |
| BILE ESCULIN AZIDE ISO 7899 APAT IRSA 29:2003 | Isolation of <i>Enterococcus</i> spp. | 1832 | 20 polystyrene tubes | 270 days | 8-25 | |
| BILE ESCULIN AZIDE ISO 7899 APAT IRSA 29:2003 | Isolation of <i>Enterococcus</i> spp. | 6005 | Dehydrated 500 gr | 3 years | 8-25 | |
| BILE ESCULIN AZIDE UNI EN 15788 | Isolation of <i>Enterococcus</i> spp. | 21476 | 20 plates 90 mm | 270 days | 8-25 | |
| BILE ESCULIN AZIDE UNI EN 15788 | Isolation of <i>Enterococcus</i> spp. | 21477 | 40 plates 60 mm | 270 days | 8-25 | |
| BILE ESCULIN AZIDE BROTH | Isolation of <i>Enterococcus</i> spp. | 1152 | 10 glass tubes | 270 days | 8-25 | |
| BILE ESCULIN AZIDE BROTH | Isolation of <i>Enterococcus</i> spp. | 1852 | 20 polystyrene tubes | 270 days | 8-25 | |
| BILE ESCULIN AZIDE BROTH | Isolation of <i>Enterococcus</i> spp. | 6273 | Dehydrated 500 gr | 3 years | 8-25 | |
| BILE SALT N° 3 | For use in microbiology | 6405 | 500 gr | 3 years | 8-25 | |
| BISMUTH SULFITE AGAR (WILSON BLAIR) USP | For the isolation of <i>Salmonella</i> spp. especially for <i>S. typhi</i> | 4 | 20 plates 90 mm | 180 days | 4-8 | |
| BISMUTH SULFITE AGAR (WILSON BLAIR) USP | For the isolation of <i>Salmonella</i> spp. especially for <i>S. typhi</i> | 20787 | 40 plates 60 mm | 180 days | 4-8 | |
| BISMUTH SULFITE AGAR (WILSON BLAIR) USP | For the isolation of <i>Salmonella</i> spp. especially for <i>S. typhi</i> | 6192 | Dehydrated 500 gr | 3 years | 8-25 | |
| BLOOD AGAR BASE | For the determination haemolytic activity of fastidious microorganisms | 6007 | Dehydrated 500 gr To use with "BLOOD, SHEEP DEBRIFINATED STERILE" | 3 years | 8-25 | |
| BLOOD AGAR BASE + NALIDIXIC ACID | For the differentiation of hemolytic activity of <i>Streptococci</i> and <i>L. monocytogenes</i> | 6274 | Dehydrated 500 gr Supplement required: "BLOOD, SHEEP DEBRIFINATED STERILE" | 3 years | 8-25 | |
| BLOOD AGAR ISO 7932 AND 11290 AMD 2004 BLOOD AGAR ISO 7932 AND 11290 AMD 2004 | For the determination of <i>Bacillus cereus</i> and <i>Listeria</i> spp. | 20343 | 20 plates 90 mm | 90 days | 4-8 | |

| | | | | | | |
|-------------------------------------|--|-------|---|----------|------|--|
| BLOOD, HORSE AGAR | General use media and study of haemolytic reactions | 1022 | 20 plates 90 mm | 90 days | 4-8 | |
| BLOOD, COLUMBIA | General use media and study of haemolytic reactions | 1023 | 20 plates 90 mm | 90 days | 4-8 | |
| BLOOD, COLUMBIA 5% (EP) | General use media and study of haemolytic reactions | 20407 | 20 plates 90 mm | 90 days | 4-8 | |
| BLOOD, COLUMBIA CNA | For the isolation of Staphylococci and Streptococci | 1048 | 20 plates 90 mm | 90 days | 4-8 | |
| BLOOD, COLUMBIA CNA MOD. | For the isolation and the differentiation of Streptococci according to hemolysis | 1024 | 20 plates 90 mm | 90 days | 4-8 | |
| BORDET GENGOU AGAR BASE | For the culture of Bordetella pertussis | 6008 | Dehydrated 500 gr Supplement required: "BLOOD, SHEEP DEBRIFINATED STERILE" | 3 years | 8-25 | |
| BORDETELLA SUPPLEMENT | To make selective Bordet Gengou agar base | 6302 | 10 vials x 500 ml | 3 years | 4-8 | |
| BORDETELLA PERTUSSIS | For the detection of Bordetella pertussis | 1036 | 20 plates 90 mm | 90 days | 4-8 | |
| BRAIN HEARTH AGAR | For the culture of fastidious microorganisms | 1005 | 20 plates 90 mm | 270 days | 8-25 | |
| BRAIN HEARTH AGAR | For the culture of fastidious microorganisms | 1105 | 10 tubes slant | 270 days | 8-25 | |
| BRAIN HEARTH AGAR | For the culture of fastidious microorganisms | 1805 | 20 polystyrene tubes slant | 270 days | 8-25 | |
| BRAIN HEARTH AGAR | For the culture of fastidious microorganisms | 1205 | 4 bottles 100 ml | 270 days | 8-25 | |
| BRAIN HEARTH AGAR | For the culture of fastidious microorganisms | 6009 | Dehydrated 500 gr | 3 years | 8-25 | |
| BRAIN HEARTHBROTH ISO 6888-1 | For the culture of fastidious microorganisms | 1155 | 10 glass tubes | 270 days | 8-25 | |
| BRAIN HEARTHBROTH ISO 6888-1 | For the culture of fastidious microorganisms | 20014 | 10 glass tubes 5 ml | 270 days | 8-25 | |
| BRAIN HEARTHBROTH ISO 6888-1 | For the culture of fastidious microorganisms | 20213 | 20 polystyrene tubes 2 ml | 270 days | 8-25 | |
| BRAIN HEARTHBROTH ISO 6888-1 | For the culture of fastidious microorganisms | 1855 | 20 polystyrene tubes | 270 days | 8-25 | |
| BRAIN HEARTHBROTH ISO 6888-1 | For the culture of fastidious microorganisms | 1255 | 4 bottles 100 ml | 270 days | 8-25 | |
| BRAIN HEARTHBROTH ISO 6888-1 | For the culture of fastidious microorganisms | 20761 | 4 bottles 200 ml | 270 days | 8-25 | |
| BRAIN HEARTHBROTH ISO 6888-1 | For the culture of fastidious microorganisms | 6010 | Dehydrated 500 gr | 3 years | 8-25 | |
| BRILLIANT GREEN (EP) | For the isolation of Salmonella spp. except S. typhi | 1082 | 20 plates 90 mm | 180 days | 4-8 | |
| BRILLIANT GREEN (EP) | For the isolation of Salmonella spp. except S. typhi | 1282 | 4 bottles 100 ml | 180 days | 4-8 | |
| BRILLIANT GREEN (EP) | For the isolation of Salmonella spp. except S. typhi | 4082 | 40 contact 55 mm | 180 days | 4-8 | |
| BRILLIANT GREEN (EP) | For the isolation of Salmonella spp. except S. typhi | 20788 | 40 plates 60 mm | 180 days | 4-8 | |
| BRILLIANT GREEN (EP) | For the isolation of Salmonella spp. except S. typhi | 6245 | Dehydrated 500 gr | 3 years | 8-25 | |
| BRILLIANT GREEN MODIFIED (ISO 6579) | For the isolation of Salmonella spp. except S. typhi | 1056 | 20 plates 90 mm | 180 days | 4-8 | |
| BRILLIANT GREEN MODIFIED (ISO 6579) | For the isolation of Salmonella spp. except S. typhi | 20276 | plates10 plates 150 mm | 180 days | 4-8 | |
| BRILLIANT GREEN MODIFIED (ISO 6579) | For the isolation of Salmonella spp. except S. typhi | 1256 | 4 bottles 100 ml | 180 days | 4-8 | |
| BRILLIANT GREEN MODIFIED (ISO 6579) | For the isolation of Salmonella spp. except S. typhi | 6833 | Dehydrated 500 gr | 3 years | 8-25 | |

| | | | | | | |
|---|---|-------|---|----------|------|--|
| BRILLIANT GREEN BILE BROTH 2% | For the detection of Coliforms in water and food | 1153 | 10 glass tubes C | 270 days | 8-25 | |
| BRILLIANT GREEN BILE BROTH 2% | For the detection of Coliforms in water and food | 1253 | 4 bottles 100 ml | 270 days | 8-25 | |
| BRILLIANT GREEN BILE BROTH 2% | For the detection of Coliforms in water and food | 6224 | Dehydrated 500 gr | 3 years | 8-25 | |
| BRILLIANT GREEN BILE BROTH 2% 2X | For the detection of Coliforms in water and food | 20170 | 10 glass tubes C | 270 days | 8-25 | |
| BRILLIANT GREEN SULFA (USDA-FSIS) | For the isolation of <i>Salmonella</i> spp. except <i>S. typhi</i> | 1035 | 20 plates 90 mm | 180 days | 4-8 | |
| BRILLIANT GREEN SULFA (USDA-FSIS) | For the isolation of <i>Salmonella</i> spp. except <i>S. typhi</i> | 1235 | 4 bottles 100 ml | 180 days | 4-8 | |
| BRILLIANT GREEN SULFA (USDA-FSIS) | For the isolation of <i>Salmonella</i> spp. except <i>S. typhi</i> | 6892 | Dehydrated 500 gr | 3 years | 8-25 | |
| BRUCELLA SELECTIVE | For the isolation of Brucella spp. | 20194 | 20 plates 90 mm | 90 days | 4-8 | |
| BRUCELLA AGAR BASE | For the isolation of Brucella spp. | 6014 | Dehydrated 500 gr Or with "BLOOD, SHEEP DEBRIFIINATED STERILE" | 3 years | 8-25 | |
| BRUCELLA AGAR BASE – REQUIRED SUPPLEMENT | For the isolation of Brucella spp. | 6301 | 10 vials x 500 ml / BRUCELLA SUPPLEMENT | 3 years | 4-8 | |
| BRUCELLA BROTH ISO 10272 | For the culture of Brucella spp. and <i>Campylobacter</i> spp. | 20294 | 10 glass tubes | 180 days | 4-8 | |
| BRUCELLA BROTH ISO 10272 | For the culture of Brucella spp. and <i>Campylobacter</i> spp. | 6015 | Dehydrated 500 gr | 3 years | 8-25 | |
| BURKHOLDERIA CEPACIA | For the isolation of Burkholderia cepacia | 20211 | 20 plates 90 mm | 180 days | 4-8 | |
| BURKHOLDERIA CEPACIA | For the isolation of Burkholderia cepacia | 20398 | 40 plates 60 mm | 180 days | 4-8 | |
| BURKHOLDERIA CEPACIA AGAR BASE | For the isolation of Burkholderia cepacia | 20421 | 4 bottles 100 ml | 3 years | 8-25 | |
| BURKHOLDERIA CEPACIA AGAR BASE | For the isolation of Burkholderia cepacia | 6285 | Dehydrated 500 gr | 180 days | 8-25 | |
| BURKHOLDERIA CEPACIA AGAR BASE – REQUIRED SUPPLEMENT | | 6332 | 10 vials x 500 ml / BURKHOLDERIA CEPACIA SUPPLEMENT | 3 years | 4-8 | |
| CALCIUM CASEINATE AGAR | For the detection of proteolytic microorganisms in food | 6286 | Dehydrated 500 gr | 3 years | 4-8 | |
| CAMPYLOBACTER BLOOD (BLASER WANG) | For the isolation of <i>Campylobacter</i> spp. | 1058 | 20 plates 90 mm | 90 days | 4-8 | |
| CAMPYLOBACTER COLUMBIA BLOOD ISO 10272 | For the culture and for the confirmation of <i>Campylobacter</i> spp. | 20293 | 20 plates 90 mm | 90 days | 4-8 | |
| CAMPYLOBACTER AGAR BASE BLOOD FREE | For the isolation of <i>Campylobacter</i> spp. | 6288 | Dehydrated 500 gr | 3 years | 8-25 | |
| CAMPYLOBACTER AGAR BASE BLOOD FREE – REQUIRED SUPPLEMENT | | 6333 | 10 vials x 500 ml / CAMPYLOBACTER BLOOD FREE SUPPLEMENT | 3 years | 4-8 | |
| CAMPYLOBACTER (CCDA BOLTON) | For the isolation of <i>Campylobacter</i> spp. | 1041 | 20 plates 90 mm | 180 days | 4-8 | |
| CAMPYLOBACTER MCCD AGAR BASE ISO 10272-1:2006 | For the isolation of <i>Campylobacter</i> spp. | 20292 | 20 plates 90 mm | 180 days | 4-8 | |
| CAMPYLOBACTER MCCD AGAR ISO 10272-1:2006 | For the isolation of <i>Campylobacter</i> spp. | 6862 | Dehydrated 500 gr | 3 years | 8-25 | |
| CAMPYLOBACTER MCCD AGAR ISO 10272-1:2006 – REQUIRED SUPPLEMENT | | 6544 | 10 vials x 500 ml / CAMPYLOBACTER MCCD SUPPLEMENT | 3 years | 4-8 | |
| CAMPYLOBACTER BOLTON BROTH ISO 10272-1 | For the selective enrichment of <i>Campylobacter</i> spp. | 20296 | 4 bottles 225 ml | 180 days | 4-8 | |
| CAMPYLOBACTER BOLTON BROTH ISO 10272-1 | For the selective enrichment of <i>Campylobacter</i> spp. | 20389 | 10 glass tubes | 180 days | 4-8 | |
| CAMPYLOBACTER BOLTON BROTH ISO 10272-1 | For the selective enrichment of <i>Campylobacter</i> spp. | 20399 | 20 polystyrene tubes | 180 days | 4-8 | |
| CAMPYLOBACTER BOLTON BROTH BASE ISO 10272-1 | For the selective enrichment of <i>Campylobacter</i> spp. | 6864 | Dehydrated 500 gr | 3 years | 8-25 | |
| CAMPYLOBACTER BOLTON BROTH BASE ISO 10272-1 – REQUIRED SUPPLEMENT | | 6543 | 10 vials x 500 ml / CAMPYLOBACTER BOLTON SUPPLEMENT | 3 years | 4-8 | |

| | | | | | | |
|---|---|-------|--|----------|------|--|
| CAMPYLOBACTER BOLTON BROTH BASE ISO 10272-1 – REQUIRED SUPPLEMENT | | 1481 | 1 bottle 50 ml / BLOOD LAKED HORSE | 365 days | 4-8 | |
| CAMPYLOBACTER KARMALI | For the determination of <i>Campylobacter</i> spp. | 20697 | 20 plates 90 mm | 180 days | 4-8 | |
| CAMPYLOBACTER AGAR BASE KARMALI | For the determination of <i>Campylobacter</i> spp. | 6704 | Dehydrated 500 gr | 3 years | 8-25 | |
| CAMPYLOBACTER AGAR BASE KARMALI – REQUIRED SUPPLEMENT | | 6500 | 10 vials x 500 ml / CAMPYLOBACTER KARMALI SUPPLEMENT | 2 anni | 4-8 | |
| CAMPYLOBACTER PRESTON ISO 10272 | For the determination of <i>Campylobacter</i> spp. | 20297 | 20 plates 90 mm | 180 days | 4-8 | |
| CAMPYLOBACTER PRESTON AGAR BASE ISO 10272 | For the isolation of <i>Campylobacter</i> spp. | 6287 | Dehydrated 500 gr | 3 years | 8-25 | |
| CAMPYLOBACTER PRESTON AGAR BASE ISO 10272 – REQUIRED SUPPLEMENT | | 6391 | 10 vials x 500 ml / CAMPYLOBACTER PRESTON SUPPLEMENT | 3 years | 4-8 | |
| CAMPYLOBACTER PRESTON AGAR BASE ISO 10272 – REQUIRED SUPPLEMENT | | 1481 | 1 bottle 50 ml / BLOOD LAKED HORSE | 365 days | 4-8 | |
| CAMPYLOBACTER PRESTON BROTH ISO 10272 | For the isolation of <i>Campylobacter</i> spp. | 20181 | 4 bottles 100 ml | 180 days | 4-8 | |
| CAMPYLOBACTER PRESTON BROTH ISO 10272 | For the isolation of <i>Campylobacter</i> spp. | 20699 | 10 glass tubes 9 ml | 180 days | 4-8 | |
| CARBOHYDRATE FERMENTATION BROTH BASE ISO 11290 | For the confirmation of <i>Listeria</i> spp. | 6776 | Dehydrated 500 gr | 3 years | 8-25 | |
| CETRIMIDE AGAR BASE (EP) | For the isolation of <i>Pseudomonas aeruginosa</i> | 6079 | Dehydrated 500 gr | 3 years | 8-25 | |
| CETRIMIDE AGAR BASE (EP) – REQUIRED SUPPLEMENT | | 6316 | 1 x 1 lt / GLYCEROL | 3 years | 8-25 | |
| CHRISTENSEN CITRATE AGAR | For the differentiation of Enterobacteria according to citrate production | 6726 | Dehydrated 500 gr | 3 years | 8-25 | |
| CHRISTENSEN CITRATE AGAR MODIFIED | For the differentiation of Enterobacteria according to citrate production | 6728 | Dehydrated 500 gr | 3 years | 8-25 | |
| C.L.E.D. | For the determination of total bacteria count in urine | 1003 | 20 plates 90 mm | 270 days | 4-8 | |
| C.L.E.D. | For the determination of total bacteria count in urine | 1203 | 4 bottles 100 ml | 270 days | 4-8 | |
| C.L.E.D. | For the determination of total bacteria count in urine | 6018 | Dehydrated 500 gr | 3 years | 4-8 | |
| CLOSTRIDIUM BOTULINUM AGAR BASE | For the isolation of <i>Clostridium botulinum</i> | 6705 | Dehydrated 500 gr | 3 years | 8-25 | |
| CLOSTRIDIUM BOTULINUM AGAR BASE – REQUIRED SUPPLEMENT | | 6501 | 10 vials x 500 ml / CLOSTRIDIUM BOTULINUM SUPPLEMENT | 2 anni | 4-8 | |
| CLOSTRIDIUM PERFRINGENS AGAR BASE (M-CP) | For the determination of <i>Clostridium perfringens</i> | 6019 | Dehydrated 500 gr | 3 years | 8-25 | |
| CLOSTRIDIUM PERFRINGENS AGAR BASE (M-CP) – REQUIRED SUPPLEMENT | | 6392 | 10 vials x 500 ml / M-CP SUPPLEMENT 1 | 3 years | 4-8 | |
| CLOSTRIDIUM PERFRINGENS AGAR BASE (M-CP) – REQUIRED SUPPLEMENT | | 6393 | 10 vials x 500 ml / M-CP SUPPLEMENT 2 | 3 years | 4-8 | |
| CLOSTRIDIUM PERFRINGENS AGAR BASE (M-CP) – REQUIRED SUPPLEMENT | | 6303 | 10 vials x 500 ml / CLOSTRIDIUM PERFRINGENS SUPPLEMENT (D-CYCLOCERINE) | 3 years | 4-8 | |
| COLUMBIA AGAR BASE (EP) | For the culture of fastidious microorganisms and the study of hemolytic reactions | 1246 | 4 bottles 100 ml Supplement required: all the products present in "ANIMAL BLOOD AND DERIVATES" | 270 days | 8-25 | |
| COLUMBIA AGAR BASE (EP) | For the culture of fastidious microorganisms and the study of hemolytic reactions | 6022 | Dehydrated 500 gr Supplement required: all the products present in "ANIMAL BLOOD AND DERIVATES" | 3 years | 8-25 | |
| COLUMBIA CNA AGAR BASE | For the isolation of Gram positive Cocci | 6021 | Dehydrated 500 gr Supplement required: all the products present in "ANIMAL BLOOD AND DERIVATES" | 3 years | 8-25 | |
| COLUMBIA HORSE BLOOD OVERLAY (USDA/FSIS) | For the determination of <i>Listeria monocytogenes</i> | 1042 | 20 plates 90 mm | 30 days | 4-8 | |

| | | | | | | |
|--|---|-------|-------------------|----------|-----|--|
| CORNMEAL AGAR + TWEEN 80 1% ISO 18416 | For the culture of yeasts and fungi | 20882 | 20 plates 90 mm | 180 days | 4-8 | |
| COUNT AGAR SUGAR FREE | Total bacterial count | 20778 | 20 plates 90 mm | 270 days | 4-8 | |
| COUNT AGAR SUGAR FREE | Total bacterial count | 20162 | 4 bottles 100 ml | 270 days | 4-8 | |
| COUNT AGAR SUGAR FREE | Total bacterial count | 6808 | Dehydrated 500 gr | 3 years | 4-8 | |
| CROMALBICANS | For the determination of C. albicans | 1093 | 20 plates 90 mm | 35 days | 4-8 | |
| CROMALBICANS | For the determination of C. albicans | 6298 | Dehydrated 500 gr | 2 anni | 4-8 | |
| CHROMOGENIC C-EC II (COLIFORMS IRSA 29:2003 7030 METHOD E) LONG EXPIRATION DATE | For the simultaneous determination of Escherichia Coli and Coliformi | 20607 | 20 plates 90 mm | 60 days | 4-8 | |
| CHROMOGENIC C-EC II (COLIFORMS IRSA 29:2003 7030 METHOD E) LONG EXPIRATION DATE | For the simultaneous determination of Escherichia Coli and Coliformi | 20501 | 40 plates 60 mm | 60 days | 4-8 | |
| CHROMOGENIC C-EC II (COLIFORMS IRSA 29:2003 7030 METHOD E) LONG EXPIRATION DATE | For the simultaneous determination of Escherichia Coli and Coliformi | 20608 | 40 contact 55 mm | 60 days | 4-8 | |
| CHROMOGENIC C-EC II (COLIFORMS IRSA 29:2003 7030 METHOD E) LONG EXPIRATION DATE | For the simultaneous determination of Escherichia Coli and Coliformi | 20541 | 4 bottles 100 ml | 60 days | 4-8 | |
| CHROMOGENIC C-EC II (COLIFORMS IRSA 29:2003 7030 METHOD E) LONG EXPIRATION DATE | For the simultaneous determination of Escherichia Coli and Coliformi | 6740 | Dehydrated 500 gr | 2 anni | 4-8 | |
| CHROMOGENIC COLIFORMS IRSA 29:2003 7010 C, 7020 B, 7030 C - LONG EXPIRATION DATE | Chromogenic and fluorogenic media for the determination of Coliforms | 1095 | 20 plates 90 mm | 60 days | 4-8 | |
| CHROMOGENIC COLIFORMS IRSA 29:2003 7010 C, 7020 B, 7030 C - LONG EXPIRATION DATE | Chromogenic and fluorogenic media for the determination of Coliforms | 2295 | 40 plates 60 mm | 60 days | 4-8 | |
| CHROMOGENIC COLIFORMS IRSA 29:2003 7010 C, 7020 B, 7030 C - LONG EXPIRATION DATE | Chromogenic and fluorogenic media for the determination of Coliforms | 4095 | 40 contact 55 mm | 60 days | 4-8 | |
| CHROMOGENIC COLIFORMS IRSA 29:2003 7010 C, 7020 B, 7030 C - LONG EXPIRATION DATE | Chromogenic and fluorogenic media for the determination of Coliforms | 1295 | 4 bottles 100 ml | 60 days | 4-8 | |
| CHROMOGENIC COLIFORMS IRSA 29:2003 7010 C, 7020 B, 7030 C - LONG EXPIRATION DATE | Chromogenic and fluorogenic media for the determination of Coliforms | 6778 | Dehydrated 500 gr | 2 anni | 4-8 | |
| CHROMOGENIC COLIFORMS (CCA) ISO 9308-1:2014 | For the simultaneous determination of E. Coli and Coliforms in water and food | 21389 | 20 plates 90 mm | 60 days | 4-8 | |
| CHROMOGENIC COLIFORMS (CCA) ISO 9308-1:2014 | For the simultaneous determination of E. Coli and Coliforms in water and food | 21390 | 40 plates 60 mm | 60 days | 4-8 | |
| CHROMOGENIC COLIFORMS (CCA) ISO 9308-1:2014 | For the simultaneous determination of E. Coli and Coliforms in water and food | 21446 | 4 bottles 100 ml | 60 days | 4-8 | |
| CHROMOGENIC COLIFORMS (CCA) ISO 9308-1:2014 | For the simultaneous determination of E. Coli and Coliforms in water and food | 6922 | Dehydrated 500 gr | 2 anni | 4-8 | |
| CHROMOGENIC E. COLI-COLIFORMS - LONG EXPIRATION DATE | For the simultaneous determination of E. Coli and Coliforms in water and food | 1097 | 20 plates 90 mm | 60 days | 4-8 | |
| CHROMOGENIC E. COLI-COLIFORMS - LONG EXPIRATION DATE | For the simultaneous determination of E. Coli and Coliforms in water and food | 2297 | 40 plates 60 mm | 60 days | 4-8 | |

| | | | | | |
|---|--|-------|---|----------|------|
| | | | | | |
| CHROMOGENIC E. COLI-COLIFORMS - LONG EXPIRATION DATE | For the simultaneous determination of E. Coli and Coliforms in water and food | 4097 | 40 contact 55 mm | 60 days | 4-8 |
| CHROMOGENIC E. COLI-COLIFORMS - LONG EXPIRATION DATE | For the simultaneous determination of E. Coli and Coliforms in water and food | 1297 | 4 bottles 100 ml | 60 days | 4-8 |
| CHROMOGENIC E. COLI-COLIFORMS - LONG EXPIRATION DATE | For the simultaneous determination of E. Coli and Coliforms in water and food | 20138 | 10 tubes 15 ml | 60 days | 4-8 |
| CHROMOGENIC E. COLI-COLIFORMS - LONG EXPIRATION DATE | For the simultaneous determination of E. Coli and Coliforms in water and food | 6232 | Dehydrated 500 gr | 2 anni | 4-8 |
| CHROMOGENIC E. COLI X-GLUC. IRSA 29.2003 7030 METHOD D - LONG EXPIRATION DATE | For the determination of E. Coli in water and food | 1099 | 20 plates 90 mm | 150 days | 4-8 |
| CHROMOGENIC E. COLI X-GLUC. IRSA 29.2003 7030 METHOD D - LONG EXPIRATION DATE | For the determination of E. Coli in water and food | 2299 | 40 plates 60 mm | 150 days | 4-8 |
| CHROMOGENIC E. COLI X-GLUC. IRSA 29.2003 7030 METHOD D - LONG EXPIRATION DATE | For the determination of E. Coli in water and food | 4099 | 40 contact 55 mm | 150 days | 4-8 |
| CHROMOGENIC E. COLI X-GLUC. IRSA 29.2003 7030 METHOD D - LONG EXPIRATION DATE | For the determination of E. Coli in water and food | 6682 | Dehydrated 500 gr | 2 anni | 4-8 |
| CHROMOGENIC E. COLI TBX LONG EXPIRATION DATE ISO 16649 | For the determination of E. Coli | 1090 | 20 plates 90 mm | 150 days | 4-8 |
| CHROMOGENIC E. COLI TBX LONG EXPIRATION DATE ISO 16649 | For the determination of E. Coli | 2290 | 40 plates 60 mm | 150 days | 4-8 |
| CHROMOGENIC E. COLI TBX LONG EXPIRATION DATE ISO 16649 | For the determination of E. Coli | 4090 | 40 contact 55 mm | 150 days | 4-8 |
| CHROMOGENIC E. COLI TBX LONG EXPIRATION DATE ISO 16649 | For the determination of E. Coli | 1290 | 4 bottles 100 ml | 150 days | 4-8 |
| CHROMOGENIC E. COLI TBX LONG EXPIRATION DATE ISO 16649 | For the determination of E. Coli | 20733 | 4 bottles 200 ml | 150 days | 4-8 |
| CHROMOGENIC E. COLI TBX LONG EXPIRATION DATE ISO 16649 | For the determination of E. Coli | 20157 | 10 glass tubes 15 ml | 150 days | 4-8 |
| CHROMOGENIC E. COLI TBX AGAR BASE ISO 16649 | For the determination of E. Coli | 6177 | Dehydrated 500 gr | 2 anni | 4-8 |
| CHROMOGENIC E. COLI TBX AGAR BASE ISO 16649 – REQUIRED SUPPLEMENT | | 12202 | 1 x 100 ml / DMSO (Dimethyl sulfoxid) | 3 anni | 8-25 |
| CHROMOGENIC LISTERIA LONG EXPIRATION DATE ISO 11290:2004 | For the isolation of Listeria spp. and presumptive identifications of L. monocytogenes | 1051 | 20 plates 90 mm | 120 days | 4-8 |
| CHROMOGENIC LISTERIA LONG EXPIRATION DATE ISO 11290:2004 | For the isolation of Listeria spp. and presumptive identifications of L. monocytogenes | 1351 | 10 plates 150 mm | 120 days | 4-8 |
| CHROMOGENIC LISTERIA LONG EXPIRATION DATE ISO 11290:2004 | For the isolation of Listeria spp. and presumptive identifications of L. monocytogenes | 4051 | 40 contact 55 mm | 120 days | 4-8 |
| CHROMOGENIC LISTERIA AGAR BASE ISO 11290:2004 | For the isolation of Listeria spp. and presumptive identifications of L. monocytogenes | 6181 | Dehydrated 500 gr | 2 anni | 4-8 |
| CHROMOGENIC LISTERIA AGAR BASE ISO 11290:2004 – REQUIRED SUPPLEMENT | | 6326 | 10 vials x 500 ml / CHROMOGENIC LISTERIA SUPPLEMENT | 3 years | 4-8 |
| CHROMOGENIC LISTERIA AGAR BASE ISO 11290:2004 – REQUIRED SUPPLEMENT | | 6335 | 10 vials x 500 ml / LISTERIA LIPASI C SUPPLEMENT | 3 years | 4-8 |
| CHROMOGENIC SALMONELLA | For the isolation of Salmonella spp. | 1096 | 20 plates 90 mm | 35 days | 4-8 |
| CHROMOGENIC SALMONELLA | For the isolation of Salmonella spp. | 2296 | 40 plates 60 mm | 35 days | 4-8 |

| | | | | | | |
|--|--|-------|-----------------------|----------|------|--|
| CHROMOGENIC SALMONELLA | For the isolation of Salmonella spp. | 4096 | 40 contact 55 mm | 35 days | 4-8 | |
| CHROMOGENIC SALMONELLA | For the isolation of Salmonella spp. | 6212 | Dehydrated 500 gr | 2 years | 4-8 | |
| CHROMOGENIC URINE LONG EXPIRATION DATE | For the count and immediate identification of urinary pathogens | 1092 | 20 plates 90 mm | 120 days | 4-8 | |
| CHROMOGENIC URINE LONG EXPIRATION DATE | For the count and immediate identification of urinary pathogens | 6660 | Dehydrated 500 gr | 2 anni | 4-8 | |
| DE-NEUTRALIZING AGAR | For the neutralization and testing of antiseptics and disinfectants | 20953 | 20 plates 90 mm | 180 days | 4-8 | |
| DE-NEUTRALIZING BROTH ISO 18416 | For the neutralization and testing of antiseptics and disinfectants | 20227 | 4 bottles 250 ml | 180 days | 4-8 | |
| DE-NEUTRALIZING BROTH ISO 18416 | For the neutralization and testing of antiseptics and disinfectants | 20609 | 4 bottles 100 ml | 180 days | 4-8 | |
| DE-NEUTRALIZING BROTH ISO 18416 | For the neutralization and testing of antiseptics and disinfectants | 20610 | 10 glass tubes | 180 days | 4-8 | |
| DE-NEUTRALIZING BROTH ISO 18416 | For the neutralization and testing of antiseptics and disinfectants | 21216 | 20 polystyrene tubes | 180 days | 4-8 | |
| DE-NEUTRALIZING BROTH BASE ISO 18416 | For the neutralization and testing of antiseptics and disinfectants | 6908 | Dehydrated 500 gr | 3 years | 8-25 | |
| DE-NEUTRALIZING BROTH BASE ISO 18416 – REQUIRED SUPPLEMENT | | 20004 | 1 x 100 ml / TWEEN 80 | - | 4-8 | |
| DERMATOPHYTES | Isolation of Dermatophytes | 1094 | 20 plates 90 mm | 180 days | 4-8 | |
| DERMATOPHYTES | Isolation of Dermatophytes | 2294 | 40 plates 60 mm | 180 days | 4-8 | |
| DERMATOPHYTES | Isolation of Dermatophytes | 3294 | 10 tubi | 180 days | 4-8 | |
| DEOXYCHOLATE | For the isolation and the enumeration of Coliforms | 1006 | 20 plates 90 mm | 210 days | 8-25 | |
| DEOXYCHOLATE | For the isolation and the enumeration of Coliforms | 6025 | Dehydrated 500 gr | 3 years | 8-25 | |
| DEOXYCHOLATE CITRATE (EP) | For the isolation of Enterobacteriaceae, especially Salmonella spp. and Shigella spp. | 1057 | 20 plates 90 mm | 210 days | 8-25 | |
| DEOXYCHOLATE CITRATE (EP) | For the isolation of Enterobacteriaceae, especially Salmonella spp. and Shigella spp. | 6026 | Dehydrated 500 gr | 3 years | 8-25 | |
| DEOXYCHOLATE LACTOSE | For the isolation and the enumeration of Coliforms | 20139 | 20 plates 90 mm | 210 days | 8-25 | |
| DEOXYCHOLATE LACTOSE | For the isolation and the enumeration of Coliforms | 6027 | Dehydrated 500 gr | 3 years | 8-25 | |
| DEXTROSE | For use in microbiology | 6406 | 500 gr | 3 years | 8-25 | |
| DEXTROSE AGAR | Total bacterial count | 6028 | Dehydrated 500 gr | 3 years | 8-25 | |
| DEXTROSE TRYPTONE AGAR | For the culture of thermophilic microorganisms | 20284 | 20 plates 90 mm | 180 days | 8-25 | |
| DEXTROSE TRYPTONE AGAR | For the culture of thermophilic microorganisms | 20350 | 4 bottles 100 ml | 180 days | 8-25 | |
| DEXTROSE TRYPTONE AGAR | For the culture of thermophilic microorganisms | 6228 | Dehydrated 500 gr | 3 years | 8-25 | |
| DEXTROSE TRYPTONE BROTH | For the culture of thermophilic microorganisms | 20750 | 4 bottles 100 ml | 180 days | 8-25 | |
| DEXTROSE TRYPTONE BROTH | For the culture of thermophilic microorganisms | 6779 | Dehydrated 500 gr | 3 years | 8-25 | |
| DG 18 ISO 21527 | For the count of yeasts and fungi in the food and feed with aw less or equal than 0,95 | 20617 | 20 plates 90 mm | 180 days | 4-8 | |
| DG 18 ISO 21527 | For the count of yeasts and fungi in the food and feed with aw less or equal than 0,95 | 20754 | 4 bottles 100 ml | 180 days | 4-8 | |

| | | | | | | |
|---|---|--------|---|----------|------|--|
| DG 18 ISO 21527 | For the count of yeasts and fungi in the food and feed with aw less or equal than 0,95 | 21278 | 10 tubes 15 ml | 180 days | 4-5 | |
| DG 18 AGAR BASE ISO 21527 | For the count of yeasts and fungi in the food and feed with aw less or equal than 0,95 | 6781 | Dehydrated 500 gr | 3 years | 4-8 | |
| DG 18 AGAR BASE ISO 21527 – REQUIRED SUPPLEMENT | | 6316 | 1 x 1 lt / GLYCEROL | 3 years | 4-8 | |
| EC BROTH ISO 7251 | For the determination of Coliforms | 1116 A | 10 glass tubes C | 180 days | 4-8 | |
| EC BROTH ISO 7251 | For the determination of Coliforms | 20312 | 4 bottles 100 ml | 180 days | 4-8 | |
| EC BROTH ISO 7251 | For the determination of Coliforms | 6246 | Dehydrated 500 gr | 3 years | 8-25 | |
| EC BROTH + MUG | For the determination of Coliforms | 20686 | 4 bottles 100 ml | 150 days | 4-8 | |
| EDWARD MEDIUM | For the isolation of Streptococcus agalactiae and other streptococci, etiological agents of bovine mastitis | 20260 | 20 plates 90 mm | 90 days | 4-8 | |
| EDWARD MEDIUM AGAR BASE | For the isolation of Streptococcus agalactiae and other streptococci, etiological agents of bovine mastitis | 6665 | Dehydrated 500 gr Supplement required: "BLOOD, SHEEP DEBRIFINATED STERILE" | 3 years | 8-25 | |
| EE MOSSEL BROTH (EP) | For the selective enrichment of Enterobacteriaceae, especially Salmonella and Coliforms | 20167 | 10 glass tubes C | 210 days | 4-8 | |
| EE MOSSEL BROTH (EP) | For the selective enrichment of Enterobacteriaceae, especially Salmonella and Coliforms | 20702 | 10 glass tubes 2X C | 210 days | 4-8 | |
| EE MOSSEL BROTH (EP) | For the selective enrichment of Enterobacteriaceae, especially Salmonella and Coliforms | 123 | 4 bottles 100 ml | 210 days | 4-8 | |
| EE MOSSEL BROTH (EP) | For the selective enrichment of Enterobacteriaceae, especially Salmonella and Coliforms | 21567 | 4 bottles 100 ml TP | 210 days | 4-8 | |
| EE MOSSEL BROTH (EP) | For the selective enrichment of Enterobacteriaceae, especially Salmonella and Coliforms | 6207 | Dehydrated 500 gr | 3 years | 8-25 | |
| EE MOSSEL BROTH ISO 7402 E 8523 | For the selective enrichment of Enterobacteriaceae, especially Salmonella and Coliforms | 20969 | 10 glass tubes C | 210 days | 4-8 | |
| EE MOSSEL BROTH ISO 7402 E 8523 | For the selective enrichment of Enterobacteriaceae, especially Salmonella and Coliforms | 20970 | 10 glass tubes 2X C | 210 days | 4-8 | |
| EE MOSSEL BROTH ISO 7402 E 8523 | For the selective enrichment of Enterobacteriaceae, especially Salmonella and Coliforms | 20971 | 4 bottles 100 ml | 210 days | 4-8 | |
| EE MOSSEL BROTH ISO 7402 E 8523 | For the selective enrichment of Enterobacteriaceae, especially Salmonella and Coliforms | 6913 | Dehydrated 500 gr | 3 years | 8-25 | |
| EMB (EOSIN METHYLENE BLUE) | For the isolation and differentiation of Coliforms from other Enterobacteriaceae | 1007 A | 20 plates 90 mm | 210 days | 8-25 | |
| EMB (EOSIN METHYLENE BLUE) | For the isolation and differentiation of Coliforms from other Enterobacteriaceae | 1207 A | 4 bottles 100 ml | 210 days | 8-25 | |
| EMB (EOSIN METHYLENE BLUE) | For the isolation and differentiation of Coliforms from other Enterobacteriaceae | 6031 | Dehydrated 500 gr | 3 years | 8-25 | |

| | | | | | | |
|--|---|-------|-------------------------------|----------|------|--|
| EMB (EOSIN METHYLENE BLUE) LEVINE | For the isolation and differentiation of Coliforms from other Enterobacteriaceae | 1007 | 20 plates 90 mm | 210 days | 4-8 | |
| EMB (EOSIN METHYLENE BLUE) LEVINE | For the isolation and differentiation of Coliforms from other Enterobacteriaceae | 1207 | 4 bottles 100 ml | 210 days | 4-8 | |
| EMB (EOSIN METHYLENE BLUE) LEVINE | For the isolation and differentiation of Coliforms from other Enterobacteriaceae | 6032 | Dehydrated 500 gr | 3 years | 8-25 | |
| ENRICHED CHOCOLATE | For the growth of Haemophilus spp. and Neisseria spp. | 1001 | 20 plates 90 mm | 270 days | 4-8 | |
| ENRICHED CHOCOLATE + BACITRACIN | For the isolation of Hemophilus spp. | 1002 | 20 plates 90 mm | 210 days | 4-8 | |
| ESCLINE IRON AGAR (E.I.A.) | For the confirmation of Enterococcus spp. | 2200 | 40 plates 60 mm | 180 days | 4-8 | |
| ETHYLVIOLET AZIDE BROTH (E.V.A.) | For the confirmation of Enterococcus spp. | 1151 | 10 glass tubes | 180 days | 8-25 | |
| ETHYLVIOLET AZIDE BROTH (E.V.A.) | For the confirmation of Enterococcus spp. | 6035 | Dehydrated 500 gr | 3 years | 8-25 | |
| EUGON LT 100 AGAR | For the eugenic growth of most microorganisms | 20568 | 4 bottles 100 ml | 180 days | 4-8 | |
| EUGON LT 100 AGAR BASE | For the eugenic growth of most microorganisms | 6975 | Dehydrated 500 gr | 3 years | 4-8 | |
| EUGON LT 100 AGAR BASE – REQUIRED SUPPLEMENT | | 6563 | 1 bottle 100 ml / OCTOXINOL 9 | 3 years | 4-8 | |
| EUGON LT 100 AGAR BASE – REQUIRED SUPPLEMENT | | 20004 | 1 bottle 100 ml / TWEEN 80 | - | 4-8 | |
| EUGON LT 100 BROTH ISO 18416 E 21149 | For the enrichment of mesophilic microorganisms in the analysis of cosmetics | 20567 | 4 bottles 100 ml | 180 days | 4-8 | |
| EUGON LT 100 BROTH ISO 18416 E 21149 | For the enrichment of mesophilic microorganisms in the analysis of cosmetics | 20869 | 10 glass tubes 9 ml | 180 days | 4-8 | |
| EUGON LT 100 BROTH ISO 18416 E 21149 | For the enrichment of mesophilic microorganisms in the analysis of cosmetics | 21209 | 4 bottles 1000 ml | 180 days | 4-8 | |
| EUGON LT 100 BROTH ISO 18416 E 21149 | For the enrichment of mesophilic microorganisms in the analysis of cosmetics | 21210 | 2 bags 3 lt | 180 days | 4-8 | |
| EUGON LT 100 BROTH BASE ISO 18416 E 21149 | For the enrichment of mesophilic microorganisms in the analysis of cosmetics | 6945 | Dehydrated 500 gr | 3 years | 4-8 | |
| EUGON LT 100 BROTH BASE ISO 18416 E 21149 – REQUIRED | | 6563 | 1 bottle 100 ml / OCTOXINOL 9 | 3 years | 4-8 | |
| EUGON LT 100 BROTH BASE ISO 18416 E 21149 BROTH BASE ISO 18416 E 21149 – REQUIRED SUPPLEMENT | | 20004 | 1 bottle 100 ml / TWEEN 80 | - | 4-8 | |
| FLUID A (EP, USP) | To aid the rinsing of the membrane filter apparatus when performing sterility testing | 20133 | 4 bottles 200 ml | 180 days | 4-20 | |
| FLUID A (EP, USP) | To aid the rinsing of the membrane filter apparatus when performing sterility testing | 130 A | 4 bottles 500 ml TP | 180 days | 4-20 | |
| FLUID A (EP, USP) | To aid the rinsing of the membrane filter apparatus when performing sterility testing | 20091 | 4 bottles 100 ml TP | 180 days | 4-20 | |
| FLUID A (EP, USP) | To aid the rinsing of the membrane filter apparatus when performing sterility testing | 20530 | 4 bottles 100 ml | 180 days | 4-20 | |
| FLUID A (EP, USP) | To aid the rinsing of the membrane filter apparatus when performing sterility testing | 20498 | 4 bottles 500 ml | 180 days | 4-20 | |
| FLUID A (EP, USP) | To aid the rinsing of the membrane filter apparatus when performing sterility testing | 20499 | 4 bottles 1000 ml | 180 days | 4-20 | |
| FLUID A (EP, USP) | To aid the rinsing of the membrane filter apparatus when performing sterility testing | 131 | 4 bottles 1000 ml TP | 180 days | 4-20 | |

| | | | | | | |
|---------------------------------|---|-------|----------------------|----------|------|--|
| | when performing sterility testing | | | | | |
| FLUID A (EP, USP) | To aid the rinsing of the membrane filter apparatus when performing sterility testing | 20827 | 4 bottles 300 ml TP | 180 days | 4-20 | |
| FLUID D (EP, USP) | To enable the rinsing of products containing lecithin or oil | 20090 | 4 bottles 100 ml TP | 180 days | 4-20 | |
| FLUID D (EP, USP) | To enable the rinsing of products containing lecithin or oil | 20531 | 4 bottles 100 ml | 180 days | 4-20 | |
| FLUID D (EP, USP) | To enable the rinsing of products containing lecithin or oil | 21358 | 4 bottles 1000 ml TP | 180 days | 4-20 | |
| FLUID K (EP, USP) | To aid the rinsing of the membrane filter apparatus when performing sterility testing | 20807 | 4 bottles 1000 ml | 180 days | 4-20 | |
| FLUID K (EP, USP) | To aid the rinsing of the membrane filter apparatus when performing sterility testing | 21359 | 4 bottles 1000 ml TP | 180 days | 4-20 | |
| GARDNERELLA SELECTIVE | For the isolation of Gardnerella vaginalis | 1040 | 20 plates 90 mm | 90 days | 4-8 | |
| GELATIN LACTOSE ISO 7937 | For the confirmation of Clostridium perfringens | 20405 | 10 glass tubes | 180 days | 4-8 | |
| GELATIN LACTOSE ISO 7937 | For the confirmation of Clostridium perfringens | 6296 | Dehydrated 500 gr | 3 years | 8-25 | |
| GELATIN, NUTRITIVE | For total bacteria count in dairy products | 1053 | 20 plates 90 mm | 210 days | 4-8 | |
| GELATIN, NUTRITIVE | For total bacteria count in dairy products | 1171 | 10 glass tubes | 210 days | 4-8 | |
| GELATIN, NUTRITIVE | For total bacteria count in dairy products | 6612 | Dehydrated 500 gr | 3 years | 4-8 | |
| GELISATO AGAR | For the study of gelatinase activity | 20131 | 10 tubes vetro 14 ml | 270 days | 4-8 | |
| GELISATO AGAR | For the study of gelatinase activity | 20868 | 4 bottles 100 ml | 270 days | 4-8 | |
| GIOLITTI CANTONI BROTH ISO 5944 | For the determination of S. aureus in food samples | 6297 | Dehydrated 500 gr | 3 years | 8-25 | |
| GLYCEROL BROTH | For the maintenance and the preservation of the bacterial strains | 20426 | 20 cryotubes 2 ml | 270 days | 4-8 | |
| GLYCEROL BROTH | For the maintenance and the preservation of the bacterial strains | 20415 | 4 bottles 100 ml | 270 days | 4-8 | |
| GLYCEROL BROTH | For the maintenance and the preservation of the bacterial strains | 20782 | 10 tubes 9 ml | 270 days | 4-8 | |
| GLUCOSE BROTH | For the study of the fermentation of glucose | 1180 | 10 glass tubes C | 270 days | 8-25 | |
| HEART INFUSION | For use in microbiology | 6432 | Dehydrated 500 gr | 3 years | 8-25 | |
| HEKTOEN ENTERIC AGAR | For the isolation and the differentiation of Gram-negative enteric bacteria | 1008 | 20 plates 90 mm | 210 days | 8-25 | |
| HEKTOEN ENTERIC AGAR | For the isolation and the differentiation of Gram-negative enteric bacteria | 1208 | 4 bottles 100 ml | 210 days | 8-25 | |
| HEKTOEN ENTERIC AGAR | For the isolation and the differentiation of Gram-negative enteric bacteria | 6040 | Dehydrated 500 gr | 3 years | 8-25 | |
| HELICOBACTER PYLORI (DENT) | For the isolation of H. pylori | 1079 | 20 plates 90 mm | 180 days | 4-8 | |
| IRON SULPHITE AGAR ISO 15213 | For the count of the anaerobic sulphite reducing bacteria | 20351 | 4 bottles 100 ml | 180 days | 4-8 | |
| IRON SULPHITE AGAR ISO 15213 | For the count of the anaerobic sulphite reducing bacteria | 6911 | Dehydrated 500 gr | 3 years | 8-25 | |
| KF STREPTOCOCCUS | For the isolation of Enterococcus spp. | 1071 | 20 plates 90 mm | 180 days | 4-8 | |
| KF STREPTOCOCCUS | For the isolation of Enterococcus spp. | 2271 | 40 plates 60 mm | 180 days | 4-8 | |
| KS STREPTOCOCCUS AGAR BASE | For the isolation of Enterococcus spp. | 6042 | Dehydrated 500 gr | 3 years | 8-25 | |

| | | | | | | |
|--|---|-------|---|----------|------|--|
| KS STREPTOCOCCUS AGAR BASE – REQUIRED SUPPLEMENT | | 6531 | 10 vials x 500 ml / TTC 1 % SUPPLEMENT | 3 years | 4-8 | |
| KING A AGAR BASE USP | For the identification of <i>Pseudomonas aeruginosa</i> | 6850 | Dehydrated 500 gr | 3 years | 8-25 | |
| KING A AGAR BASE USP – REQUIRED SUPPLEMENT | | 6316 | 1 x 1 lt / GLYCEROL | 3 years | 8-25 | |
| KING B (PSEUDOMONAS F) ISO 16266 | For the identification of <i>Pseudomonas aeruginosa</i> | 3004 | 20 plates 90 mm | 240 days | 8-25 | |
| KING B (PSEUDOMONAS F) ISO 16266 | For the identification of <i>Pseudomonas aeruginosa</i> | 20680 | 40 plates 60 mm | 240 days | 8-25 | |
| KING B (PSEUDOMONAS F) ISO 16266 | For the identification of <i>Pseudomonas aeruginosa</i> | 20023 | 10 tubes 5 ml slant | 240 days | 8-25 | |
| KING B (PSEUDOMONAS F) ISO 16266 | For the identification of <i>Pseudomonas aeruginosa</i> | 20285 | 20 polystyrene tubes slant | 240 days | 8-25 | |
| KING B (PSEUDOMONAS F) ISO 16266 | For the identification of <i>Pseudomonas aeruginosa</i> | 20564 | 4 bottles 100 ml | 240 days | 8-25 | |
| KING B (PSEUDOMONAS F) AGAR BASE ISO 16266 | For the identification of <i>Pseudomonas aeruginosa</i> | 6602 | Dehydrated 500 gr | 3 years | 8-25 | |
| KING B (PSEUDOMONAS F) AGAR BASE ISO 16266 – REQUIRED SUPPLEMENT | | 6316 | 1 x 1 lt / GLYCEROL | 3 years | 8-25 | |
| KING B MEDIUM AGAR BASE USP | For the identification of <i>Pseudomonas aeruginosa</i> | 6205 | Dehydrated 500 gr | 3 years | 8-25 | |
| KING B MEDIUM AGAR BASE USP – REQUIRED SUPPLEMENT | | 6316 | 1 x 1 lt / GLYCEROL | 3 years | 8-25 | |
| KLIGER IRON AGAR | For the differentiation of Gram-negative enterobacteria | 20783 | 20 plates 90 mm | 270 days | 8-25 | |
| KLIGER IRON AGAR | For the differentiation of Gram-negative enterobacteria | 1136 | 10 glass tubes slant | 270 days | 8-25 | |
| KLIGER IRON AGAR | For the differentiation of Gram-negative enterobacteria | 1836 | 20 polystyrene tubes slant | 270 days | 8-25 | |
| KLIGER IRON AGAR | For the differentiation of Gram-negative enterobacteria | 6043 | Dehydrated 500 gr | 3 years | 8-25 | |
| KLIGER IRON AGAR ISO 10273 | For the differentiation of Gram-negative enterobacteria | 20704 | 10 glass tubes slant | 270 days | 8-25 | |
| KLIGER IRON AGAR ISO 10273 | For the differentiation of Gram-negative enterobacteria | 6604 | Dehydrated 500 gr | 3 years | 8-25 | |
| KLIGER IRON AGAR APAT MAN. 20/2003 METODO 3 | For the differentiation of Gram-negative enterobacteria | 21052 | 10 glass tubes slant | 270 days | 8-25 | |
| KLIGER IRON AGAR APAT MAN. 20/2003 METODO 3 | For the differentiation of Gram-negative enterobacteria | 21053 | 20 polystyrene tubes slant | 270 days | 8-25 | |
| LACTOSE | For use in microbiology | 6408 | 500 gr | 3 years | 8-25 | |
| LACTOSE BROTH (EP) QUAD. ISSN 0390 days- 6239:1983 | For the detection of Coliforms and <i>Salmonella</i> spp. | 1158 | 10 glass tubes C | 270 days | 8-25 | |
| LACTOSE BROTH (EP) QUAD. ISSN 0390 days- 6239:1983 | For the detection of Coliforms and <i>Salmonella</i> spp. | 20659 | 10 tubes 9 ml C | 270 days | 8-25 | |
| LACTOSE BROTH (EP) QUAD. ISSN 0390 days- 6239:1983 | For the detection of Coliforms and <i>Salmonella</i> spp. | 1258 | 4 bottles 100 ml | 270 days | 8-25 | |
| LACTOSE BROTH (EP) QUAD. ISSN 0390 days- 6239:1983 | For the detection of Coliforms and <i>Salmonella</i> spp. | 6044 | Dehydrated 500 gr | 3 years | 8-25 | |
| LACTOSE BROTH 2X | For the detection of Coliforms | 1159 | 10 glass tubes C | 270 days | 8-25 | |
| LACTOSE SULPHITE BROTH ISO 7937 | For the isolation of <i>Clostridium perfringens</i> | 121 | 4 bottles 100 ml | 90 days | 4-8 | |
| LACTOSE SULPHITE BROTH ISO 7937 | For the isolation of <i>Clostridium perfringens</i> | 20310 | 10 glass tubes | 90 days | 4-8 | |
| LACTOSE SULPHITE BROTH ISO 7937 | For the isolation of <i>Clostridium perfringens</i> | 20627 | 10 glass tubes 5 ml | 90 days | 4-8 | |
| LACTOSE SULPHITE BROTH ISO 7937 | For the isolation of <i>Clostridium perfringens</i> | 20705 | 10 glass tubes 9 ml | 90 days | 4-8 | |
| LACTOSE SULPHITE BROTH ISO 7937 | For the isolation of <i>Clostridium perfringens</i> | 20881 | 10 glass tubes 8 ml | 90 days | 4-8 | |
| LACTOSE SULPHITE BROTH BASE ISO 7937 | For the isolation of <i>Clostridium perfringens</i> | 6197 | Dehydrated 500 gr | 3 years | 8-25 | |

| | | | | | | |
|--|---|-------|---|----------|------|--|
| LACTOSE SULPHITE BROTH BASE ISO 7937 – REQUIRED SUPPLEMENT | | 6560 | 5 + 5 fiale x 500 ml / LACTOSE SULPHITE BROTH SUPPLEMENT | 3 years | 8-25 | |
| LAURIL SULFATE AGAR | For the isolation and the enumeration of Coliforms | 6046 | Dehydrated 500 gr | 3 years | 8-25 | |
| LAURIL SULFATE BROTH (LTB) | For the determination of Coliforms in water | 1134 | 10 glass tubes C | 180 days | 8-25 | |
| LAURIL SULFATE BROTH (LTB) | For the determination of Coliforms in water | 20257 | 4 bottles 100 ml | 180 days | 8-25 | |
| LAURIL SULFATE BROTH (LTB) | For the determination of Coliforms in water | 6045 | Dehydrated 500 gr | 3 years | 8-25 | |
| LAURIL SULFATE BROTH 2X | For the determination of Coliforms in water | 20615 | 10 glass tubes C | 180 days | 8-25 | |
| LAURIL SULFATE BROTH MODIFIED ISO 22964 | For the selective enrichment of Enterobacter sakazakii | 20191 | 10 glass tubes C | 60 days | 4-8 | |
| LAURIL SULFATE BROTH MODIFIED ISO 22964 | For the selective enrichment of Enterobacter sakazakii | 20708 | 4 bottles 225 ml | 60 days | 4-8 | |
| LAURIL SULFATE BROTH MODIFIED 2X ISO 22964 | For the determination of Coliforms in water | 20707 | 10 glass tubes C | 60 days | 4-8 | |
| LB AGAR (LENNOX) | For the studies of molecular genetics with E. Coli | 6278 | Dehydrated 500 gr | 3 years | 8-25 | |
| LB BROTH (LENNOX) | For the studies of molecular genetics with E. Coli | 6279 | Dehydrated 500 gr | 3 years | 8-25 | |
| LEGIONELLA SELECTIVE GVPC ISO 11731 | For the isolation of Legionella spp. | 1077 | 20 plates 90 mm | 180 days | 4-8 | |
| LEGIONELLA SELECTIVE GVPC ISO 11731 | For the isolation of Legionella spp. | 4077 | 40 contact 55 mm | 180 days | 4-8 | |
| LEGIONELLA SELECTIVE GVPC ISO 11731 | For the isolation of Legionella spp. | 2277 | 40 plates 60 mm | 180 days | 4-8 | |
| LEGIONELLA SELECTIVE GVPC ISO 11731 | For the isolation of Legionella spp. | 20200 | 40 contact s/m | 180 days | 4-8 | |
| LEGIONELLA (BCYE) ISO 11731 | For the culture of Legionella spp. | 20500 | 20 plates 90 mm | 180 days | 4-8 | |
| LEGIONELLA (BCYE) ISO 11731 | For the culture of Legionella spp. | 20613 | 40 plates 60 mm | 180 days | 4-8 | |
| LEGIONELLA (BCYE) ISO 11731 | For the culture of Legionella spp. | 20413 | 40 contact 55 mm | 180 days | 4-8 | |
| LEGIONELLA (BCYE WITHOUT CYSTEINE) ISO 11731 | For the confirmation of Legionella spp. | 20281 | 20 plates 90 mm | 180 days | 4-8 | |
| LEGIONELLA (BCYE WITHOUT CYSTEINE) ISO 11731 | For the confirmation of Legionella spp. | 20611 | 40 plates 60 mm | 180 days | 4-8 | |
| LEGIONELLA (BCYE WITHOUT CYSTEINE) ISO 11731 | For the confirmation of Legionella spp. | 20618 | 40 contact 55 mm | 180 days | 4-8 | |
| LEGIONELLA AGAR BASE ISO 11731 | For the isolation of Legionella spp. | 6650 | Dehydrated 500 gr | 3 years | 8-25 | |
| LEGIONELLA AGAR BASE ISO 11731 – REQUIRED SUPPLEMENT | | 6382 | 10 vials x 100 ml / LEGIONELLA CYE GROWTH SUPPLEMENT | 3 years | 8-25 | |
| LEGIONELLA AGAR BASE ISO 11731 – REQUIRED SUPPLEMENT | | 6395 | 10 vials x 500 ml / LEGIONELLA GVPC SUPPLEMENT | 3 years | 8-25 | |
| LETHEEN AGAR BASE MODIFIED | For the determination bactericidal activity of ammonium quaternary salt | 6605 | Dehydrated 500 gr | 3 years | 8-25 | |
| LETHEEN AGAR BASE MODIFIED – REQUIRED SUPPLEMENT | | 20004 | 1 x 100 ml / TWEEN 80 | - | 4-8 | |
| LETHEEN BROTH BASE MODIFIED ISO 18416 | For the determination bactericidal activity of ammonium quaternary salt | 6606 | Dehydrated 500 gr | 3 years | 8-25 | |
| LETHEEN BROTH BASE MODIFIED ISO 18416 – REQUIRED SUPPLEMENT | | 20004 | 1 x 100 ml / TWEEN 80 | - | 4-8 | |
| LETHEEN MODIFIED BROTH ISO 18416 | For the determination bactericidal activity of ammonium quaternary salt | 20182 | 10 glass tubes 9 ml | 180 days | 8-25 | |
| LETHEEN MODIFIED BROTH ISO 18416 | For the determination bactericidal activity of ammonium quaternary salt | 20601 | 10 glass tubes | 180 days | 8-25 | |
| LETHEEN MODIFIED BROTH ISO 18416 | For the determination bactericidal activity of ammonium quaternary salt | 13 | 4 bottles 100 ml | 180 days | 8-25 | |
| LISTERIA BUFFERED ENRICHMENT BROTH | For the isolation of Listeria spp. | 21015 | 10 glass tubes | 120 days | 4-8 | |

| | | | | | | |
|--|------------------------------------|---------|--|----------|------|--|
| LISTERIA BUFFERED ENRICHMENT BROTH | For the isolation of Listeria spp. | 21015,5 | 50 glass tubes | 120 days | 4-8 | |
| LISTERIA BUFFERED ENRICHMENT BROTH | For the isolation of Listeria spp. | 1259 A | 4 bottles 225 ml | 120 days | 4-8 | |
| LISTERIA BUFFERED ENRICHMENT BROTH | For the isolation of Listeria spp. | 20987 | 2 bags 3 lt | 120 days | 4-8 | |
| LISTERIA BUFFERED ENRICHMENT BROTH | For the isolation of Listeria spp. | 20988 | 2 bags 5 lt | 120 days | 4-8 | |
| LISTERIA BUFFERED ENRICHMENT BROTH | For the isolation of Listeria spp. | 6231 | Dehydrated 500 gr | 3 years | 8-25 | |
| LISTERIA ENRICHMENT BROTH FDA | For the isolation of Listeria spp. | 6783 | Dehydrated 500 gr | 3 years | 8-25 | |
| LISTERIA FRASER BROTH ISO 11290 | For the isolation of Listeria spp. | 1102 | 10 glass tubes | 120 days | 4-8 | |
| LISTERIA FRASER BROTH ISO 11290 | For the isolation of Listeria spp. | 1202 | 4 bottles 100 ml | 120 days | 4-8 | |
| LISTERIA FRASER BROTH ISO 11290 | For the isolation of Listeria spp. | 20210 | 4 bottles 225 ml | 120 days | 4-8 | |
| LISTERIA FRASER BROTH ISO 11290 | For the isolation of Listeria spp. | 20846 | 2 bags 3 lt | 120 days | 4-8 | |
| LISTERIA FRASER BROTH ISO 11290 | For the isolation of Listeria spp. | 21000 | 2 bags 5 lt | 120 days | 4-8 | |
| LISTERIA FRASER BROTH HALF CONCENTRATION ISO 11290 | For the isolation of Listeria spp. | 1102 A | 10 glass tubes | 120 days | 4-8 | |
| LISTERIA FRASER BROTH HALF CONCENTRATION ISO 11290 | For the isolation of Listeria spp. | 1202 A | 4 bottles 100 ml | 120 days | 4-8 | |
| LISTERIA FRASER BROTH HALF CONCENTRATION ISO 11290 | For the isolation of Listeria spp. | 20204 | 4 bottles 225 ml | 120 days | 4-8 | |
| LISTERIA FRASER BROTH HALF CONCENTRATION ISO 11290 | For the isolation of Listeria spp. | 20820 | 4 bottles 250 ml | 120 days | 4-8 | |
| LISTERIA FRASER BROTH HALF CONCENTRATION ISO 11290 | For the isolation of Listeria spp. | 20900 | 2 bags 3 lt | 120 days | 4-8 | |
| LISTERIA FRASER BROTH HALF CONCENTRATION ISO 11290 | For the isolation of Listeria spp. | 20901 | 2 bags 5 lt | 120 days | 4-8 | |
| LISTERIA FRASER HALF CONCENTRATION WITHOUT FERRIC AMMONIUM CITRATE | For the isolation of Listeria spp. | 21325 | 4 bottles 225 ml | 120 days | 4-8 | |
| LISTERIA FRASER HALF CONCENTRATION WITHOUT FERRIC AMMONIUM CITRATE | For the isolation of Listeria spp. | 21328 | 10 tubes 10 ml | 120 days | 4-8 | |
| LISTERIA FRASER HALF CONCENTRATION WITHOUT FERRIC AMMONIUM CITRATE | For the isolation of Listeria spp. | 21384 | 2 bags 3 lt | 120 days | 4-8 | |
| LISTERIA FRASER HALF CONCENTRATION WITHOUT FERRIC AMMONIUM CITRATE | For the isolation of Listeria spp. | 21385 | 2 bags 5 lt | 120 days | 4-8 | |
| LISTERIA FRASER BROTH BASE ISO 11290 | For the isolation of Listeria spp. | 6047 | Dehydrated 500 gr | 3 years | 8-25 | |
| LISTERIA FRASER BROTH BASE ISO 11290 – TO USE WITH | | 6305 | 2 x 5 vials x 500 ml / FRASER LISTERIA SELECTIVE SUPPLEMENT | 3 years | 4-8 | |
| LISTERIA FRASER BROTH BASE ISO 11290 – TO USE WITH | | 6306 | 2 x 5 vials x 500 ml / HALF FRASER LISTERIA SELECTIVE SUPPLEMENT | 3 years | 4-8 | |
| LISTERIA OXFORD ISO 11290 | For the isolation of Listeria spp. | 1063 | 20 plates 90 mm | 180 days | 4-8 | |
| LISTERIA OXFORD ISO 11290 | For the isolation of Listeria spp. | 4063 | 40 contact 55 mm | 180 days | 4-8 | |
| LISTERIA OXFORD AGAR BASE ISO 11290 | For the isolation of Listeria spp. | 6071 | Dehydrated 500 gr | 3 years | 8-25 | |
| LISTERIA OXFORD AGAR BASE ISO 11290 – REQUIRED SUPPLEMENT | | 6307 | 10 vials x 500 ml / OXFORD LISTERIA SELECTIVE SUPPLEMENT | 3 years | 4-8 | |
| LISTERIA PALCAM ISO 11290 | For the isolation of Listeria spp. | 1062 | 20 plates 90 mm | 180 days | 4-8 | |
| LISTERIA PALCAM ISO 11290 | For the isolation of Listeria spp. | 4062 | 40 contact 55 mm | 180 days | 4-8 | |
| LISTERIA PALCAM ISO 11290 | For the isolation of Listeria spp. | 1897 | 20 polystyrene tubes 5 ml slant | 180 days | 4-8 | |

| | | | | | | |
|---|---|--------|--|----------|------|--|
| LISTERIA PALCAM ISO 11290 | For the isolation of Listeria spp. | 1197 | 10 glass tubes slant | 180 days | 4-8 | |
| LISTERIA PALCAM AGAR BASE ISO 11290 | For the isolation of Listeria spp. | 6199 | Dehydrated 500 gr | 3 years | 8-25 | |
| LISTERIA PALCAM AGAR BASE ISO 11290 – REQUIRED SUPPLEMENT | | 6308 | 10 vials x 500 ml / PALCAM LISTERIA SELECTIVE SUPPLEMENT | 3 years | 4-8 | |
| LISTERIA SELECTIVE FAST BROTH | For the selective enrichment of Listeria in 18 hours!! | 21382 | 4 bottles 225 ml | 180 days | 4-8 | |
| LISTERIA SELECTIVE FAST BROTH | For the selective enrichment of Listeria in 18 hours!! | 21333 | 2 bags 3 lt | 180 days | 4-8 | |
| LISTERIA SELECTIVE FAST BROTH | For the selective enrichment of Listeria in 18 hours!! | 21334 | 2 bags 5 lt | 180 days | 4-8 | |
| LISTERIA SELECTIVE FAST BROTH WITHOUT FERRIC AMMONIUM CITRATE | For the selective enrichment of Listeria in 18 hours!! | 21335 | 4 bottles 225 ml | 180 days | 4-8 | |
| LISTERIA SELECTIVE FAST BROTH WITHOUT FERRIC AMMONIUM CITRATE | For the selective enrichment of Listeria in 18 hours!! | 21355 | 2 bags 3 lt | 180 days | 4-8 | |
| LISTERIA SELECTIVE FAST BROTH WITHOUT FERRIC AMMONIUM CITRATE | For the selective enrichment of Listeria in 18 hours!! | 21356 | 2 bags 5 lt | 180 days | 4-8 | |
| LISTERIA SELECTIVE FAST BROTH BASE | For the selective enrichment of Listeria in 18 hours!! | 6971 | Dehydrated 500 gr | 3 years | 8-25 | |
| LISTERIA SELECTIVE FAST BROTH BASE – REQUIRED SUPPLEMENT | | 6583 | 10 vials x 500 ml / FERRIC AMMONIUM CITRATE | 3 years | 8-25 | |
| LISTERIA UVM1 BROTH (USDA-FSIS) | For the isolation of Listeria spp. | 1121 | 10 glass tubes | 120 days | 8-25 | |
| LISTERIA UVM1 BROTH (USDA-FSIS) | For the isolation of Listeria spp. | 1221 | 4 bottles 100 ml | 120 days | 8-25 | |
| LISTERIA UVM1 BROTH (USDA-FSIS) | For the isolation of Listeria spp. | 1221 A | 4 bottles 225 ml | 120 days | 8-25 | |
| LISTERIA UVM1 BROTH (USDA-FSIS) | For the isolation of Listeria spp. | 20893 | 2 bags 3 lt | 120 days | 8-25 | |
| LISTERIA UVM1 BROTH (USDA-FSIS) | For the isolation of Listeria spp. | 20897 | 2 bags 5 lt | 120 days | 8-25 | |
| LISTERIA UVM1 BROTH (USDA-FSIS) | For the isolation of Listeria spp. | 6638 | Dehydrated 500 gr | 3 anni | 8-25 | |
| LISTERIA UVM2 BROTH (USDA-FSIS) | For the isolation of Listeria spp. | 1122 | 10 glass tubes | 120 days | 8-25 | |
| LISTERIA UVM2 BROTH (USDA-FSIS) | For the isolation of Listeria spp. | 1222 | 4 bottles 100 ml | 120 days | 8-25 | |
| LISTERIA UVM2 BROTH (USDA-FSIS) | For the isolation of Listeria spp. | 20894 | 2 bags 3 lt | 120 days | 8-25 | |
| LISTERIA UVM2 BROTH (USDA-FSIS) | For the isolation of Listeria spp. | 20898 | 2 bags 5 lt | 120 days | 8-25 | |
| LISTERIA UVM2 BROTH (USDA-FSIS) | For the isolation of Listeria spp. | 6639 | Dehydrated 500 gr | 3 years | 8-25 | |
| LOEFFLER | For the culture of <i>Corynebacterium</i> spp. | 1181 | 10 glass tubes slant | 180 days | 4-8 | |
| LPT DILUTION BROTH | For microbiological analysis of cosmetics | 20452 | 10 glass tubes 9 ml | 210 days | 8-25 | |
| LPT DILUTION BROTH | For microbiological analysis of cosmetics | 1231 | 4 bottles 100 ml | 210 days | 8-25 | |
| LPT DILUTION BROTH | For microbiological analysis of cosmetics | 20523 | 4 bottles 200 ml | 210 days | 8-25 | |
| LPT DILUTION BROTH | For microbiological analysis of cosmetics | 20552 | 4 bottles 500 ml | 210 days | 8-25 | |
| LPT DILUTION BROTH BASE | For microbiological analysis of cosmetics | 6226 | Dehydrated 500 gr | 3 years | 8-25 | |
| LPT DILUTION BROTH BASE – REQUIRED SUPPLEMENT | | 20004 | 1 x 100 ml / TWEEN 80 | - | 4-8 | |
| LURIA AGAR MILLER'S LB AGAR | For <i>E.coli</i> molecular genetic studies | 6282 | Dehydrated 500 gr | 3 years | 8-25 | |
| LURIA BROTH MILLER'S LB BROTH | For the studies of molecular genetics with <i>E. Coli</i> | 6281 | Dehydrated 500 gr | 3 years | 8-25 | |
| LURIA AGAR MILLER'S MODIFICATION | For <i>E.coli</i> molecular genetic studies | 6280 | Dehydrated 500 gr | 3 years | 8-25 | |
| LURIA BROTH MILLER'S MODIFICATION | For <i>E.coli</i> molecular genetic studies | 6256 | Dehydrated 500 gr | 3 years | 8-25 | |
| LYSINE DECARBOXYLASE BROTH ISO 6579 | For the confirmation of <i>Salmonella</i> spp. | 20150 | 10 tubes 5 ml | 210 days | 8-25 | |
| LYSINE DECARBOXYLASE BROTH ISO 6579 | For the confirmation of <i>Salmonella</i> spp. | 20151 | 4 bottles 100 ml | 210 days | 8-25 | |
| LYSINE DECARBOXYLASE BROTH ISO 6579 | For the confirmation of <i>Salmonella</i> spp. | 6608 | Dehydrated 500 gr | 3 years | 8-25 | |

| | | | | | | |
|--|--|--------|---|----------|------|--|
| M 17 AGAR (APHA) | For the culture and the enumeration of lactic Streptococci | 21317 | 20 plates 90 mm | 180 days | 8-25 | |
| M 17 AGAR (APHA) | For the culture and the enumeration of lactic Streptococci | 20718 | 10 glass tubes 14 ml | 180 days | 8-25 | |
| M 17 AGAR (APHA) | For the culture and the enumeration of lactic Streptococci | 21318 | 4 bottles 100 ml | 180 days | 8-25 | |
| M 17 AGAR (APHA) | For the culture and the enumeration of lactic Streptococci | 6050 | Dehydrated 500 gr | 3 years | 8-25 | |
| MAC CONKEY AGAR (EP) | For the isolation and the differentiation of Enterobacteria | 1011 | 20 plates 90 mm | 240 days | 8-25 | |
| MAC CONKEY AGAR (EP) | For the isolation and the differentiation of Enterobacteria | 2211 | 40 plates 60 mm | 240 days | 8-25 | |
| MAC CONKEY AGAR (EP) | For the isolation and the differentiation of Enterobacteria | 4011 | 40 contact 55 mm | 240 days | 8-25 | |
| MAC CONKEY AGAR (EP) | For the isolation and the differentiation of Enterobacteria | 2511 | 40 contact s/m40 contact | 240 days | 8-25 | |
| MAC CONKEY AGAR (EP) | For the isolation and the differentiation of Enterobacteria | 1211 | 4 bottles 100 ml | 240 days | 8-25 | |
| MAC CONKEY AGAR (EP) | For the isolation and the differentiation of Enterobacteria | 6051 | Dehydrated 500 gr | 3 years | 8-25 | |
| MAC CONKEY + MUG | Selective and differential medium for the determination of Coliforms and for the direct determination of E. coli | 20113 | 20 plates 90 mm | 90 days | 4-8 | |
| MAC CONKEY + SORBITOL ISO 16654 | For the identification of Escherichia coli O 157: H7 | 1011 A | 20 plates 90 mm | 240 days | 8-25 | |
| MAC CONKEY + SORBITOL ISO 16654 | For the identification of Escherichia coli O 157: H7 | 6053 | Dehydrated 500 gr | 3 years | 8-25 | |
| MAC CONKEY + SORBITOL ISO 16654 – REQUIRED SUPPLEMENT | | 6507 | 10 vials x 500 ml / CEFIXIME + TELLURITE SUPPLEMENT | 3 years | 4-8 | |
| MAC CONKEY + SORBITOL + POTASSIUM TELLURITE CEFIXIME ISO 16654 | For the detection of Escherichia coli O 157 | 20508 | 20 plates 90 mm | 180 days | 4-8 | |
| MAC CONKEY BROTH (EP) | For the determination of Coliforms in water and food | 1137 | 10 glass tubes C | 210 days | 8-25 | |
| MAC CONKEY BROTH (EP) | For the determination of Coliforms in water and food | 122 | 4 bottles 100 ml | 210 days | 8-25 | |
| MAC CONKEY BROTH (EP) | For the determination of Coliforms in water and food | 122,4 | 40 bottles 100 ml | 210 days | 8-25 | |
| MAC CONKEY BROTH (EP) | For the determination of Coliforms in water and food | 21568 | 4 bottles 100 ml TP | 210 days | 8-25 | |
| MAC CONKEY BROTH (EP) | For the determination of Coliforms in water and food | 20637 | 4 bottles 500 ml | 210 days | 8-25 | |
| MAC CONKEY BROTH (EP) | For the determination of Coliforms in water and food | 20638 | 4 bottles 1000 ml | 210 days | 8-25 | |
| MAC CONKEY BROTH (EP) | For the determination of Coliforms in water and food | 6055 | Dehydrated 500 gr | 3 years | 8-25 | |
| MAC CONKEY BROTH 2X (EP) | For the determination of Coliforms in water and food | 1137 A | 10 glass tubes C | 210 days | 8-25 | |
| MALASSEZIA | For the growth of M. furfur and M. pachydermatis | 1075 | 20 plates 90 mm | 180 days | 4-8 | |
| MALASSEZIA | For the growth of M. furfur and M. pachydermatis | 2275 | 40 plates 60 mm | 180 days | 4-8 | |
| MALT EXTRACT | For use in microbiology | 6409 | Dehydrated 500 gr | 3 years | 8-25 | |
| MALT EXTRACT AGAR | For the growth of yeasts and fungi | 20022 | 20 plates 90 mm | 180 days | 8-25 | |
| MALT EXTRACT AGAR | For the growth of yeasts and fungi | 20489 | 4 bottles 100 ml | 180 days | 8-25 | |
| MALT EXTRACT AGAR | For the growth of yeasts and fungi | 20740 | 40 contact s/m | 180 days | 8-25 | |
| MALT EXTRACT AGAR | For the growth of yeasts and fungi | 20021 | 40 contact 55 mm | 180 days | 8-25 | |
| MALT EXTRACT AGAR | For the growth of yeasts and fungi | 20691 | 20 contact 90 mm s/m | 180 days | 8-25 | |

| | | | | | | |
|---|--|--------|---------------------------------------|----------|------|--|
| MALT EXTRACT AGAR | For the growth of yeasts and fungi | 6056 | Dehydrated 500 gr | 3 years | 8-25 | |
| MALT EXTRACT BROTH | For the growth of yeasts and fungi | 6255 | Dehydrated 500 gr | 3 years | 8-25 | |
| MALTOSE | For use in microbiology | 6420 | Dehydrated 500 gr | 3 years | 8-25 | |
| MANNITOL SALT (EP) | For the isolation and the differentiation of Staphylococci | 1019 | 20 plates 90 mm | 210 days | 8-25 | |
| MANNITOL SALT (EP) | For the isolation and the differentiation of Staphylococci | 2219 | 40 plates 60 mm | 210 days | 8-25 | |
| MANNITOL SALT (EP) | For the isolation and the differentiation of Staphylococci | 4019 | 40 contact 55 mm | 210 days | 8-25 | |
| MANNITOL SALT (EP) | For the isolation and the differentiation of Staphylococci | 2519 | 40 contact s/m | 210 days | 8-25 | |
| MANNITOL SALT (EP) | For the isolation and the differentiation of Staphylococci | 1119 | 10 glass tubes slant | 210 days | 8-25 | |
| MANNITOL SALT (EP) | For the isolation and the differentiation of Staphylococci | 1219 | 4 bottles 100 ml | 210 days | 8-25 | |
| MANNITOL SALT (EP) | For the isolation and the differentiation of Staphylococci | 1219 B | 4 bottles 250 ml | 210 days | 8-25 | |
| MANNITOL SALT (EP) | For the isolation and the differentiation of Staphylococci | 6058 | Dehydrated 500 gr | 3 years | 8-25 | |
| MARINE AGAR | For the isolation and the enumeration of heterotrophic marine bacteria | 6240 | Dehydrated 500 gr | 3 years | 8-25 | |
| MARINE BROTH | For the isolation and the enumeration of heterotrophic marine bacteria | 6241 | Dehydrated 500 gr | 3 years | 8-25 | |
| M-CP | For the isolation of Clostridium perfringens | 1055 | 20 plates 90 mm | 180 days | 4-8 | |
| M-CP | For the isolation of Clostridium perfringens | 2255 | 40 plates 60 mm | 180 days | 4-8 | |
| M-ENDO LES ISO 9308 | For the isolation and the enumeration of Coliforms by membrane filter method | 1084 | 20 plates 90 mm | 120 days | 4-8 | |
| M-ENDO LES ISO 9308 | For the isolation and the enumeration of Coliforms by membrane filter method | 2284 | 40 plates 60 mm | 120 days | 4-8 | |
| M-ENDO LES ISO 9308 | For the isolation and the enumeration of Coliforms by membrane filter method | 20558 | 4 bottles 100 ml | 120 days | 4-8 | |
| M-ENDO LES AGAR BASE ISO 9308 | For the count and the detection of Coliforms in the waters by membrane filter method | 6642 | Dehydrated 500 gr | 3 years | 8-25 | |
| M-ENDO LES AGAR BASE ISO 9308 – REQUIRED SUPPLEMENT | | 6312 | 25 gr / BASIC FUCHSIN | 1 year | 8-25 | |
| M-ENTEROCCUS (SLANETZ BARTLEY) ISO 7899 | For the isolation of Enterococcus by membrane filter method | 1098 | 20 plates 90 mm | 180 days | 4-8 | |
| M-ENTEROCCUS (SLANETZ BARTLEY) ISO 7899 | For the isolation of Enterococcus by membrane filter method | 2298 | 40 plates 60 mm | 180 days | 4-8 | |
| M-ENTEROCCUS (SLANETZ BARTLEY) ISO 7899 | For the isolation of Enterococcus by membrane filter method | 4098 | 40 contact | 180 days | 4-8 | |
| M-ENTEROCCUS (SLANETZ BARTLEY) ISO 7899 | For the isolation of Enterococcus by membrane filter method | 1298 | 4 bottles 100 ml | 180 days | 4-8 | |
| M-ENTEROCCUS (SLANETZ BARTLEY) AGAR BASE ISO 7899 | For the isolation of Enterococcus by membrane filter method | 6059 | Dehydrated 500 gr | 3 years | 8-25 | |
| M-ENTEROCCUS (SLANETZ BARTLEY) AGAR BASE ISO 7899 – REQUIRED SUPPLEMENT | | 6531 | 10 vials x 500 ml / TTC 1% Supplement | 3 years | 4-8 | |
| M-FC | For the count of Coliforms in the waters by filtering membrane method | 1091 | 20 plates 90 mm | 120 days | 4-8 | |

| | | | | | | |
|--|---|--------|---|----------|------|--|
| M-FC | For the count of Coliforms in the waters by filtering membrane method | 2291 | 40 plates 60 mm | 120 days | 4-8 | |
| M-FC (FECAL COLIFORMS AGAR BASE) | For the count of Coliforms in the waters by filtering membrane method | 6036 | Dehydrated 500 gr | 3 years | 8-25 | |
| M-FC (FECAL COLIFORMS AGAR BASE) – REQUIRED SUPPLEMENT | | 6334 | 10 vials x 500 ml / FECAL COLIFORMS SUPPLEMENT (ROSOLIC ACID) | 3 years | 8-25 | |
| MICOGRAMMA | For the sensitivity test to antimycotic | 20160 | 10 tubes 15 ml | 270 days | 8-25 | |
| MINERAL MODIFIED GLUTAMMATE AGAR ISO 16649 | For the count of Coliforms | 20377 | 20 plates 90 mm | 180 days | 4-8 | |
| MINERAL MODIFIED GLUTAMMATE AGAR ISO 16649 | For the count of Coliforms | 6693 | Dehydrated 500 gr | 3 years | 8-25 | |
| MINERAL MODIFIED GLUTAMMATE BROTH ISO 16649 | For the count of Coliforms | 20391 | 10 glass tubes C | 180 days | 4-8 | |
| MINERAL MODIFIED GLUTAMMATE BROTH ISO 16649 | For the count of Coliforms | 6692 | Dehydrated 500 gr | 3 years | 8-25 | |
| MINERAL MODIFIED GLUTAMMATE BROTH 2X ISO 16649 | For the count of Coliforms | 20392 | 10 glass tubes C | 180 days | 4-8 | |
| MOX AGAR (USDA-FSIS) | For the isolation of Listeria spp. | 1065 | 20 plates 90 mm | 180 days | 4-8 | |
| MOX AGAR BASE (USDA-FSIS) | For the isolation of Listeria spp. | 1265 | 4 bottles 100 ml | 180 days | 4-8 | |
| MOX AGAR BASE (USDA-FSIS) | For the isolation of Listeria spp. | 6856 A | Dehydrated 500 gr | 3 years | 8-25 | |
| MOX AGAR BASE (USDA-FSIS) – REQUIRED SUPPLEMENT | | 6503 | 10 vials x 500 ml / MOXALACTAM MOXALACTAM | 3 years | 4-8 | |
| MRS + TWEEN 80 | For the growth of lactobacilli | 1083 | 20 plates 90 mm | 180 days | 4-8 | |
| MRS + TWEEN 80 | For the growth of lactobacilli | 1283 | 4 bottles 100 ml | 180 days | 4-8 | |
| MRS + TWEEN 80 | For the growth of lactobacilli | 21349 | 10 tubes 15 ml | 180 days | 4-8 | |
| MRS + TWEEN 80 | For the growth of lactobacilli | 20674 | 10 glass tubes 22 ml | 180 days | 4-8 | |
| MRS AGAR BASE | For the growth of lactobacilli | 6060 | Dehydrated 500 gr | 3 years | 4-8 | |
| MRS AGAR BASE – REQUIRED SUPPLEMENT | | 20004 | 1 x 100 ml / TWEEN 80 | - | 4-8 | |
| MRS BROTH BASE | For the growth of lactobacilli | 6061 | Dehydrated 500 gr | 3 years | 4-8 | |
| MRS BROTH BASE – REQUIRED SUPPLEMENT | | 20004 | 1 x 100 ml / TWEEN 80 | - | 4-8 | |
| MRS + CIPROFLOXACIN + CLINDAMYCIN | For the growth of lactobacilli | 20438 | 20 plates 90 mm | 90 days | 4-8 | |
| MR-VP ISO 6579 | For the differentiation of Escherichia from the other Enterobacteria | 20001 | 10 glass tubes 5 ml | 270 days | 4-8 | |
| MR-VP ISO 6579 | For the differentiation of Escherichia from the other Enterobacteria | 20725 | 10 glass tubes 3 ml | 270 days | 4-8 | |
| MR-VP ISO 6579 | For the differentiation of Escherichia from the other Enterobacteria | 20467 | 4 bottles 100 ml | 270 days | 4-8 | |
| MR-VP ISO 6579 | For the differentiation of Escherichia from the other Enterobacteria | 6062 | Dehydrated 500 gr | 3 years | 8-25 | |
| MUELLER HINTON | For sensitivity test on antibiotics by Kirby-Bauer method | 1012 | 20 plates 90 mm | 270 days | 8-25 | |
| MUELLER HINTON | For sensitivity test on antibiotics by Kirby-Bauer method | 1312 | plates10 plates 150 mm | 270 days | 8-25 | |
| MUELLER HINTON | For sensitivity test on antibiotics by Kirby-Bauer method | 1112 | 10 glass tubes 22 ml | 270 days | 8-25 | |
| MUELLER HINTON | For sensitivity test on antibiotics by Kirby-Bauer method | 1212 | 4 bottles 100 ml | 270 days | 8-25 | |

| | | | | | | |
|--|--|--------|--|----------|------|--|
| MUELLER HINTON | For sensitivity test on antibiotics by Kirby-Bauer method | 6064 | Dehydrated 500 gr | 3 years | 8-25 | |
| MUELLER HINTON BROTH | For the sensitivity testing to antibiotics | 1160 | 10 glass tubes | 270 days | 8-25 | |
| MUELLER HINTON BROTH | For the sensitivity testing to antibiotics | 1860 | 20 polystyrene tubes | 270 days | 8-25 | |
| MUELLER HINTON BROTH | For the sensitivity testing to antibiotics | 1260 | 4 bottles 100 ml | 270 days | 8-25 | |
| MUELLER HINTON BROTH | For the sensitivity testing to antibiotics | 6063 | Dehydrated 500 gr | 3 years | 8-25 | |
| MUELLER HINTON + BLOOD | For the sensitivity test of fastidious organisms and for general use | 1013 | 20 plates 90 mm | 90 days | 4-8 | |
| MUELLER HINTON + BLOOD | For the sensitivity test of fastidious organisms and for general use | 1313 | 10 plates 150 mm | 90 days | 4-8 | |
| MUELLER KAUFFMAN BROTH | For the selective enrichment of Samonella spp. In foods | 1161 | 10 glass tubes | 90 days | 4-8 | |
| MUELLER KAUFFMAN BROTH | For the selective enrichment of Samonella spp. In foods | 1861 | 20 polystyrene tubes | 90 days | 4-8 | |
| MUELLER KAUFFMAN BROTH | For the selective enrichment of Samonella spp.in foods. | 1261 | 4 bottles 100 ml | 90 days | 4-8 | |
| MUELLER KAUFFMAN BROTH BASE | For the selective enrichment of Salmonella spp. in foods | 6201 | Dehydrated 500 gr | 3 years | 8-25 | |
| MUELLER KAUFFMAN BROTH BASE – REQUIRED SUPPLEMENT | For the selective enrichment of Samonella spp. In foods. | 6360,1 | 10 vials x 500 ml / IODO-IODINE SOLUTION | 90 days | 4-8 | |
| MUELLER KAUFFMAN + NOVOBIOCIN 40 mg/l ISO 6579 | For the selective enrichment of Samonella spp. In foods | 20026 | 10 glass tubes | 90 days | 4-8 | |
| MUELLER KAUFFMAN + NOVOBIOCIN 40 mg/l ISO 6579 | For the selective enrichment of Samonella spp. In foods | 20657 | 10 glass tubes 9 ml | 90 days | 4-8 | |
| MUELLER KAUFFMAN + NOVOBIOCIN 40 mg/l ISO 6579 | For the selective enrichment of Samonella spp. In foods. | 20631 | 20 polystyrene tubes | 90 days | 4-8 | |
| MUELLER KAUFFMAN + NOVOBIOCIN 40 mg/l ISO 6579 | For the selective enrichment of Samonella spp. In foods. | 20553 | 4 bottles 100 ml | 90 days | 4-8 | |
| MUELLER KAUFFMAN + NOVOBIOCIN 40 mg/l ISO 6579 | For the selective enrichment of Samonella spp. In foods. | 21360 | 4 bottles 225 ml | 90 days | 4-8 | |
| MUELLER KAUFFMAN + NOVOBIOCIN 40 mg/l ISO 6579 | For the selective enrichment of Samonella spp. In foods. | 6202 | Dehydrated 500 gr | 3 years | 8-25 | |
| MUELLER KAUFFMAN + NOVOBIOCIN 40 mg/l ISO 6579 – REQUIRED SUPPLEMENT | For the selective enrichment of Samonella spp. In foods. | 6360,1 | 10 vials x 500 ml / IODO-IODINE SOLUTION | 90 days | 4-8 | |
| MYCOLOGICAL AGAR | For the isolation of pathogenic fungi | 1088 | 20 plates 90 mm | 180 days | 4-8 | |
| MYCOLOGICAL AGAR | For the isolation of pathogenic fungi | 1288 | 4 bottles 100 ml | 180 days | 4-8 | |
| MYCOLOGICAL AGAR | For the isolation of pathogenic fungi | 6065 | Dehydrated 500 gr | 3 years | 8-25 | |
| NEISSERIA SELECTIVE | For the isolation of Neisseria spp. | 1033 | 20 plates 90 mm | 180 days | 4-8 | |
| NEUTRALIZER BUFFER ISO 18593 | To neutralize the residues of most common disinfectants from surfaces in the food sector | 20437 | 4 bottles 100 ml | 180 days | 4-8 | |
| NEUTRALIZER BUFFER ISO 18593 | To neutralize the residues of most common disinfectants from surfaces in the food sector | 20763 | 10 glass tubes | 180 days | 4-8 | |
| NSA | For the culture of Pseudomonas spp | 3009 | 20 plates 90 mm | 210 days | 8-25 | |
| NUTRIENT AGAR ISO 6579 | For general use | 1004 | 20 plates 90 mm | 270 days | 8-25 | |
| NUTRIENT AGAR ISO 6579 | For general use | 1804 | 20 polystyrene tubes slant | 270 days | 8-25 | |
| NUTRIENT AGAR ISO 6579 | For general use | 1104 | 10 glass tubes slant | 270 days | 8-25 | |
| NUTRIENT AGAR ISO 6579 | For general use | 1204 | 4 bottles 100 ml | 270 days | 8-25 | |
| NUTRIENT AGAR ISO 6579 | For general use | 20355 | 4 bottles 200 ml | 270 days | 8-25 | |
| NUTRIENT AGAR ISO 6579 | For general use | 20762 | 4 bottles 250 ml | 270 days | 8-25 | |

| | | | | | | |
|--|---|-------|-------------------------------------|----------|------|--|
| NUTRIENT AGAR ISO 6579 | For general use | 6066 | Dehydrated 500 gr | 3 years | 8-25 | |
| NUTRIENT AGAR ISO 16266 ISTISAN 96/35 | For the identification of <i>Pseudomonas aeruginosa</i> by membrane filter method | 20519 | 20 plates 90 mm | 270 days | 8-25 | |
| NUTRIENT AGAR ISO 16266 ISTISAN 96/35 | For the identification of <i>Pseudomonas aeruginosa</i> by membrane filter method | 20993 | 40 plates 60 mm | 270 days | 8-25 | |
| NUTRIENT AGAR ISO 16266 ISTISAN 96/35 | For the identification of <i>Pseudomonas aeruginosa</i> by membrane filter method | 20518 | 4 bottles 100 ml | 270 days | 8-25 | |
| NUTRIENT AGAR ISO 16266 ISTISAN 96/35 | For the identification of <i>Pseudomonas aeruginosa</i> by membrane filter method | 6794 | Dehydrated 500 gr | 3 years | 8-25 | |
| NUTRIENT AGAR WITH SODIUM CHLORIDE ISO 21528-1 | For the confirmation of Enterobacteria | 20472 | 4 bottles 100 ml | 270 days | 8-25 | |
| NUTRIENT AGAR WITH SODIUM CHLORIDE ISO 21528-1 | For the confirmation of Enterobacteria | 20719 | 20 plates 90 mm | 270 days | 8-25 | |
| NUTRIENT AGAR WITH SODIUM CHLORIDE ISO 21528-1 | For the confirmation of Enterobacteria | 20473 | 4 bottles 200 ml | 270 days | 8-25 | |
| NUTRIENT AGAR WITH SODIUM CHLORIDE ISO 21528-1 | For the confirmation of Enterobacteria | 6610 | Dehydrated 500 gr | 3 years | 8-25 | |
| NUTRIENT AGAR SEMISOLID ISO 6579 | For general use | 20354 | 4 bottles 200 ml | 270 days | 8-25 | |
| NUTRIENT AGAR SEMISOLID ISO 6579 | For general use | 20353 | 4 bottles 100 ml | 270 days | 8-25 | |
| NUTRIENT AGAR SEMISOLID ISO 6579 | For general use | 6786 | Dehydrated 500 gr | 3 years | 8-25 | |
| NUTRIENT BROTH | For general use | 1154 | 10 glass tubes | 270 days | 8-25 | |
| NUTRIENT BROTH | For general use | 1854 | 20 polystyrene tubes | 270 days | 8-25 | |
| NUTRIENT BROTH | For general use | 1254 | 4 bottles 100 ml | 270 days | 8-25 | |
| NUTRIENT BROTH | For general use | 6067 | Dehydrated 500 gr | 3 years | 8-25 | |
| OGYE ISO 7954 | For the enumeration of yeasts and molds in the food | 1073 | 20 plates 90 mm | 150 days | 4-8 | |
| OGYE AGAR BASE ISO 7954 | For the enumeration of yeasts and molds in the food | 6711 | Dehydrated 500 gr | 3 years | 8-25 | |
| OGYE AGAR BASE ISO 7954 – REQUIRED SUPPLEMENT | | 6515 | 10 vials x 500 ml / OGYE SUPPLEMENT | 3 years | 4-8 | |
| ORANGE SERUM AGAR | For the isolation of acid-tolerant pathogens in fruit juices | 20861 | 20 plates 90 mm | 180 days | 8-25 | |
| ORANGE SERUM AGAR | For the isolation of acid-tolerant pathogens in fruit juices | 20836 | 40 plates 60 mm | 180 days | 8-25 | |
| ORANGE SERUM AGAR | For the isolation of acid-tolerant pathogens in fruit juices | 6940 | Dehydrated 500 gr | 3 years | 8-25 | |
| ORANGE SERUM AGAR IFU N° 2 | For the isolation of acid-tolerant pathogens in fruit juices | 21168 | 20 plates 90 mm | 180 days | 8-25 | |
| OVERLAY MEDIUM ISO 4833 | | 20441 | 4 bottles 150 ml | 210 days | 8-25 | |
| OVERLAY MEDIUM ISO 4833 | | 20764 | 10 glass tubes 4 ml | 210 days | 8-25 | |
| PAGE SALINE ISO 11731 | For the preparation of the biofilm and sediment suspension for Legionella spp. research | 20145 | 10 glass tubes | 270 days | 4-8 | |
| PAGE SALINE ISO 11731 | For the preparation of the biofilm and sediment suspension for Legionella spp. research | 20612 | 4 bottles 100 ml | 270 days | 4-8 | |
| PAGE SALINE ISO 11731 | For the preparation of the biofilm and sediment suspension for Legionella spp. research | 20805 | 10 glass tubes 9 ml | 270 days | 4-8 | |
| PAGE SALINE ISO 11731 | For the preparation of the biofilm and sediment suspension for Legionella spp. Research | 20806 | 4 bottles 90 ml | 270 days | 4-8 | |

| | | | | | | |
|--|--|--------|----------------------|----------|------|--|
| POTATO DEXTROSE AGAR (EP) | For the enumeration of yeasts and molds | 20192 | 20 plates 90 mm | 210 days | 8-25 | |
| POTATO DEXTROSE AGAR (EP) | For the enumeration of yeasts and molds | 20751 | 40 plates 60 mm | 210 days | 8-25 | |
| POTATO DEXTROSE AGAR (EP) | For the enumeration of yeasts and molds | 20306 | 4 bottles 100 ml | 210 days | 8-25 | |
| POTATO DEXTROSE AGAR (EP) | For the enumeration of yeasts and molds | 6078 | Dehydrated 500 gr | 3 years | 8-25 | |
| POTATO DEXTROSE BROTH | For the culture of yeasts and molds | 6244 | Dehydrated 500 gr | 3 years | 8-25 | |
| PCA (APHA) PLATE COUNT AGAR ISO 4833 | For the total aerobic bacterial count in food, water, air and surfaces | 1047 | 20 plates 90 mm | 240 days | 8-25 | |
| PCA (APHA) PLATE COUNT AGAR ISO 4833 | For the total aerobic bacterial count in food, water, air and surfaces | 2247 | 40 plates 60 mm | 240 days | 8-25 | |
| PCA (APHA) PLATE COUNT AGAR ISO 4833 | For the total aerobic bacterial count in food, water, air and surfaces | 4047 | 40 contact 55 mm | 240 days | 8-25 | |
| PCA (APHA) PLATE COUNT AGAR ISO 4833 | For the total aerobic bacterial count in food, water, air and surfaces | 2547 | 40 contact s/m | 240 days | 8-25 | |
| PCA (APHA) PLATE COUNT AGAR ISO 4833 | For the total aerobic bacterial count in food, water, air and surfaces | 20248 | 20 contact 90 mm | 240 days | 8-25 | |
| PCA (APHA) PLATE COUNT AGAR ISO 4833 | For the total aerobic bacterial count in food, water, air and surfaces | 1147 | 10 glass tubes 22 ml | 240 days | 8-25 | |
| PCA (APHA) PLATE COUNT AGAR ISO 4833 | For the total aerobic bacterial count in food, water, air and surfaces | 20130 | 10 glass tubes 15 ml | 240 days | 8-25 | |
| PCA (APHA) PLATE COUNT AGAR ISO 4833 | For the total aerobic bacterial count in food, water, air and surfaces | 1247 | 4 bottles 100 ml | 240 days | 8-25 | |
| PCA (APHA) PLATE COUNT AGAR ISO 4833 | For the total aerobic bacterial count in food, water, air and surfaces | 20044 | 4 bottles 200 ml | 240 days | 8-25 | |
| PCA (APHA) PLATE COUNT AGAR ISO 4833 | For the total aerobic bacterial count in food, water, air and surfaces | 20300 | 4 bottles 250 ml | 240 days | 8-25 | |
| PCA (APHA) PLATE COUNT AGAR ISO 4833 | For the total aerobic bacterial count in food, water, air and surfaces | 20446 | 4 bottles 500 ml | 240 days | 8-25 | |
| PCA (APHA) PLATE COUNT AGAR ISO 4833 | For the total aerobic bacterial count in food, water, air and surfaces | 6077 | Dehydrated 500 gr | 3 years | 8-25 | |
| PCA + LECITHIN + TWEEN 80 | For the total aerobic bacterial count in food, water, air and surfaces | 4047 A | 40 contact 55 mm | 180 days | 4-8 | |
| PCA + LECITHIN + TWEEN 80 | For the total aerobic bacterial count in food, water, air and surfaces | 2547 A | 40 contact s/m | 180 days | 4-8 | |
| PCA + LECITHIN + TWEEN 80 | For the total aerobic bacterial count in food, water, air and surfaces | 20249 | 20 contact 90 mm | 180 days | 4-8 | |
| PCA + LECITHIN + TWEEN 80 + HISTIDINE + SODIUM THIOSULFATE | For the total aerobic bacterial count in food, water, air and surfaces | 4047 B | 40 contact 55 mm | 180 days | 4-8 | |
| PCA + TTC | For the total aerobic bacterial count in food, water, air and surfaces | 20345 | 40 plates 60 mm | 180 days | 4-8 | |
| PCA + TTC | For the total aerobic bacterial count in food, water, air and surfaces | 20721 | 40 contact 55 mm | 180 days | 4-8 | |
| PEPTONE, ACID CASEINE | For use in microbiology | 6423 | Dehydrated 500 gr | 3 years | 8-25 | |
| PEPTONE, BACTERIOLOGICAL | For use in microbiology | 6421 | Dehydrated 500 gr | 3 years | 8-25 | |
| PEPTONE, CASEINE | For use in microbiology | 6419 | Dehydrated 500 gr | 3 years | 8-25 | |
| PEPTONE, GELATIN | For use in microbiology | 6407 | Dehydrated 500 gr | 3 years | 8-25 | |
| PEPTONE, MEAT | For use in microbiology | 6410 | Dehydrated 500 gr | 3 years | 8-25 | |
| PEPTONE, PROTEOSE | For use in microbiology | 6411 | Dehydrated 500 gr | 3 years | 8-25 | |
| PEPTONE, SOY | For use in microbiology | 6414 | Dehydrated 500 gr | 3 years | 8-25 | |
| PEPTONE WATER | For the homogenization of microbiological samples | 1150 | 10 glass tubes | 270 days | 8-25 | |

| | | | | | | |
|---|---|---------|----------------------|----------|------|--|
| PEPTONE WATER | For the homogenization of microbiological samples | 20000 | 10 glass tubes 5 ml | 270 days | 8-25 | |
| PEPTONE WATER | For the homogenization of microbiological samples | 1150,5 | 50 glass tubes | 270 days | 8-25 | |
| PEPTONE WATER | For the homogenization of microbiological samples | 1850 | 20 polystyrene tubes | 270 days | 8-25 | |
| PEPTONE WATER | For the homogenization of microbiological samples | 1250 | 4 bottles 100 ml | 270 days | 8-25 | |
| PEPTONE WATER | For the homogenization of microbiological samples | 1250 Z | 4 bottles 200 ml | 270 days | 8-25 | |
| PEPTONE WATER | For the homogenization of microbiological samples | 1250 D | 4 bottles 225 ml | 270 days | 8-25 | |
| PEPTONE WATER | For the homogenization of microbiological samples | 20283 | 4 bottles 250 ml | 270 days | 8-25 | |
| PEPTONE WATER | For the homogenization of microbiological samples | 1250 A | 4 bottles 90 ml | 270 days | 8-25 | |
| PEPTONE WATER | For the homogenization of microbiological samples | 20630 | 4 bottles 500 ml | 270 days | 8-25 | |
| PEPTONE WATER | For the homogenization of microbiological samples | 20985 | 2 bags 3 lt | 270 days | 8-25 | |
| PEPTONE WATER | For the homogenization of microbiological samples | 20986 | 2 bags 5 lt | 270 days | 8-25 | |
| PEPTONE WATER | For the homogenization of microbiological samples | 6072 | Dehydrated 500 gr | 3 years | 8-25 | |
| PEPTONE WATER (MAXIMUM RECOVERY DILUENT) ISO 6887 e ISO 18416 - ISTISAN 96/35 | For the dilution of microbiological samples | 20427 | 10 glass tubes | 270 days | 8-25 | |
| PEPTONE WATER (MAXIMUM RECOVERY DILUENT) ISO 6887 e ISO 18416 - ISTISAN 96/35 | For the dilution of microbiological samples | 20428 | 10 glass tubes 9 ml | 270 days | 8-25 | |
| PEPTONE WATER (MAXIMUM RECOVERY DILUENT) ISO 6887 e ISO 18416 - ISTISAN 96/35 | For the dilution of microbiological samples | 20428,5 | 50 glass tubes 9 ml | 270 days | 8-25 | |
| PEPTONE WATER (MAXIMUM RECOVERY DILUENT) ISO 6887 e ISO 18416 - ISTISAN 96/35 | For the dilution of microbiological samples | 20427,5 | 50 glass tubes | 270 days | 8-25 | |
| PEPTONE WATER (MAXIMUM RECOVERY DILUENT) ISO 6887 e ISO 18416 - ISTISAN 96/35 | For the dilution of microbiological samples | 20207 | 4 bottles 100 ml | 270 days | 8-25 | |
| PEPTONE WATER (MAXIMUM RECOVERY DILUENT) ISO 6887 e ISO 18416 - ISTISAN 96/35 | For the dilution of microbiological samples | 20207,4 | 40 bottles 100 ml | 270 days | 8-25 | |
| PEPTONE WATER (MAXIMUM RECOVERY DILUENT) ISO 6887 e ISO 18416 - ISTISAN 96/35 | For the dilution of microbiological samples | 20193 | 4 bottles 90 ml | 270 days | 8-25 | |
| PEPTONE WATER (MAXIMUM RECOVERY DILUENT) ISO 6887 e ISO 18416 - ISTISAN 96/35 | For the dilution of microbiological samples | 20193,4 | 40 bottles 90 ml | 270 days | 8-25 | |
| PEPTONE WATER (MAXIMUM RECOVERY DILUENT) ISO 6887 e ISO 18416 - ISTISAN 96/35 | For the dilution of microbiological samples | 20339 | 4 bottles 225 ml | 270 days | 8-25 | |
| PEPTONE WATER (MAXIMUM RECOVERY DILUENT) ISO 6887 e ISO 18416 - ISTISAN 96/35 | For the dilution of microbiological samples | 20698 | 4 bottles 250 ml | 270 days | 8-25 | |
| PEPTONE WATER (MAXIMUM RECOVERY DILUENT) ISO 6887 e ISO 18416 - ISTISAN 96/35 | For the dilution of microbiological samples | 20734 | 4 bottles 200 ml | 270 days | 8-25 | |
| PEPTONE WATER (MAXIMUM RECOVERY DILUENT) ISO 6887 e ISO 18416 - ISTISAN 96/35 | For the dilution of microbiological samples | 20396 | 4 bottles 500 ml TP | 270 days | 8-25 | |
| PEPTONE WATER (MAXIMUM RECOVERY DILUENT) ISO 6887 e ISO 18416 - ISTISAN 96/35 | For the dilution of microbiological samples | 20203 | 4 bottles 500 ml | 270 days | 8-25 | |
| PEPTONE WATER (MAXIMUM RECOVERY DILUENT) ISO 6887 e ISO 18416 - ISTISAN 96/35 | For the dilution of microbiological samples | 20831 | 4 bottles 1000 ml | 270 days | 8-25 | |

| | | | | | | |
|--|--|-----------|---------------------------|----------|------|--|
| PEPTONE WATER (MAXIMUM RECOVERY DILUENT) ISO 6887 e ISO 18416 - ISTISAN 96/35 | For the dilution of microbiological samples | 20892 | 2 bags 3 lt | 270 days | 8-25 | |
| PEPTONE WATER (MAXIMUM RECOVERY DILUENT) ISO 6887 e ISO 18416 - ISTISAN 96/35 | For the dilution of microbiological samples | 20896 | 2 bags 5 lt | 270 days | 8-25 | |
| PEPTONE WATER (MAXIMUM RECOVERY DILUENT) ISO 6887 e ISO 18416 - ISTISAN 96/35 | For the dilution of microbiological samples | 6777 | Dehydrated 500 gr | 3 years | 8-25 | |
| PEPTONE WATER, ALKALINE - ISTISAN 63/35 | For the enrichment of Vibrio spp. | 1151 A | 10 glass tubes | 270 days | 8-25 | |
| PEPTONE WATER, ALKALINE - ISTISAN 63/35 | For the enrichment of Vibrio spp. | 20668 | 10 glass tubes 9 ml | 270 days | 8-25 | |
| PEPTONE WATER, ALKALINE - ISTISAN 63/35 | For the enrichment of Vibrio spp. | 1851 A | 20 polystyrene tubes | 270 days | 8-25 | |
| PEPTONE WATER, ALKALINE - ISTISAN 63/35 | For the enrichment of Vibrio spp. | 20340 | 4 bottles 250 ml | 270 days | 8-25 | |
| PEPTONE WATER, ALKALINE - ISTISAN 63/35 | For the enrichment of Vibrio spp. | 6270 | Dehydrated 500 gr | 3 years | 8-25 | |
| PEPTONE WATER, BUFFERED (USDA/FSIS) | For the dilution of food samples | 1150 B | 10 glass tubes | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED (USDA/FSIS) | For the dilution of food samples | 1150 B.50 | 50 glass tubes | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED (USDA/FSIS) | For the dilution of food samples | 20649 | 20 polystyrene tubes 9 ml | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED (USDA/FSIS) | For the dilution of food samples | 1850 B | 20 polystyrene tubes | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED (USDA/FSIS) | For the dilution of food samples | 1250 B | 4 bottles 100 ml | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED (USDA/FSIS) | For the dilution of food samples | 20394 | 4 bottles 250 ml | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED (USDA/FSIS) | For the dilution of food samples | 20735 | 4 bottles 500 ml | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED (USDA/FSIS) | For the dilution of food samples | 20520 | 4 bottles 1000 ml | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED (USDA/FSIS) | For the dilution of food samples | 6073 | Dehydrated 500 gr | 3 years | 8-25 | |
| PEPTONE WATER, BUFFERED ISO 6579 E ISO 6887 E ISO 22964 - ISTISAN 96/35 | For the pre-enrichment of Salmonella spp. (ISO 6579) and for the preparation of decimal dilutions of the sample (ISO 6887) - For the pre-enrichment of Enterobacter sakazakii in milk and its derivatives | 20656 | 10 glass tubes 9 ml | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED ISO 6579 E ISO 6887 E ISO 22964 - ISTISAN 96/35 | For the pre-enrichment of Salmonella spp. (ISO 6579) and for the preparation of decimal dilutions of the sample (ISO 6887) - For the pre-enrichment of Enterobacter sakazakii in milk and its derivatives | 20656,5 | 50 glass tubes 9 ml | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED ISO 6579 E ISO 6887 E ISO 22964 - ISTISAN 96/35 | For the pre-enrichment of Salmonella spp. (ISO 6579) and for the preparation of decimal dilutions of the sample (ISO 6887) - For the pre-enrichment of Enterobacter sakazakii in milk and its derivatives | 20731 | 4 bottles 90 ml | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED ISO 6579 E ISO 6887 E ISO 22964 - ISTISAN 96/35 | For the pre-enrichment of Salmonella spp. (ISO 6579) and for the preparation of decimal dilutions of the sample (ISO 6887) - For the pre-enrichment of Enterobacter sakazakii in milk and its derivatives | 20731,4 | 40 bottles 90 ml | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED ISO 6579 E ISO 6887 E ISO 22964 - ISTISAN 96/35 | For the pre-enrichment of Salmonella spp. (ISO 6579) and for the preparation of decimal dilutions of the sample (ISO 6887) - For the pre-enrichment of Enterobacter sakazakii in milk and its derivatives | 20732 | 4 bottles 99 ml | 270 days | 8-25 | |

| | | | | | | |
|--|--|---------|---------------------|----------|------|--|
| PEPTONE WATER, BUFFERED ISO 6579 E ISO 6887 E ISO 22964 - ISTISAN 96/35 | For the pre-enrichment of Salmonella spp. (ISO 6579) and for the preparation of decimal dilutions of the sample (ISO 6887) - For the pre-enrichment of Enterobacter sakazakii in milk and its derivatives | 20732,4 | 40 bottles 99 ml | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED ISO 6579 E ISO 6887 E ISO 22964 - ISTISAN 96/35 | For the pre-enrichment of Salmonella spp. (ISO 6579) and for the preparation of decimal dilutions of the sample (ISO 6887) - For the pre-enrichment of Enterobacter sakazakii in milk and its derivatives | 20228 | 4 bottles 250 ml | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED ISO 6579 E ISO 6887 E ISO 22964 - ISTISAN 96/35 | For the pre-enrichment of Salmonella spp. (ISO 6579) and for the preparation of decimal dilutions of the sample (ISO 6887) - For the pre-enrichment of Enterobacter sakazakii in milk and its derivatives | 20229 | 4 bottles 500 ml | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED ISO 6579 E ISO 6887 E ISO 22964 - ISTISAN 96/35 | For the pre-enrichment of Salmonella spp. (ISO 6579) and for the preparation of decimal dilutions of the sample (ISO 6887) - For the pre-enrichment of Enterobacter sakazakii in milk and its derivatives | 20396 | 4 bottles 500 ml TP | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED ISO 6579 E ISO 6887 E ISO 22964 - ISTISAN 96/35 | For the pre-enrichment of Salmonella spp. (ISO 6579) and for the preparation of decimal dilutions of the sample (ISO 6887) - For the pre-enrichment of Enterobacter sakazakii in milk and its derivatives | 20470 | 4 bottles 225 ml | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED ISO 6579 E ISO 6887 E ISO 22964 - ISTISAN 96/35 | For the pre-enrichment of Salmonella spp. (ISO 6579) and for the preparation of decimal dilutions of the sample (ISO 6887) - For the pre-enrichment of Enterobacter sakazakii in milk and its derivatives | 20471 | 4 bottles 100 ml | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED ISO 6579 E ISO 6887 E ISO 22964 - ISTISAN 96/35 | For the pre-enrichment of Salmonella spp. (ISO 6579) and for the preparation of decimal dilutions of the sample (ISO 6887) - For the pre-enrichment of Enterobacter sakazakii in milk and its derivatives | 20516 | 4 bottles 1 lt | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED ISO 6579 E ISO 6887 E ISO 22964 - ISTISAN 96/35 | For the pre-enrichment of Salmonella spp. (ISO 6579) and for the preparation of decimal dilutions of the sample (ISO 6887) - For the pre-enrichment of Enterobacter sakazakii in milk and its derivatives | 20766 | 2 bags 3 lt | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED ISO 6579 E ISO 6887 E ISO 22964 - ISTISAN 96/35 | For the pre-enrichment of Salmonella spp. (ISO 6579) and for the preparation of decimal dilutions of the sample (ISO 6887) - For the pre-enrichment of Enterobacter sakazakii in milk and its derivatives | 20899 | 2 bags 5 lt | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED ISO 6579 E ISO 6887 E ISO 22964 - ISTISAN 96/35 | For the pre-enrichment of Salmonella spp. (ISO 6579) and for the preparation of decimal dilutions of the sample (ISO 6887) - For the pre-enrichment of Enterobacter sakazakii in milk and its derivatives | 6687 | Dehydrated 500 gr | 3 years | 8-25 | |

| | | | | | | |
|--|--|---------|---------------------------------------|----------|------|--|
| PEPTONE WATER, BUFFERED (EP) | For the dilution of microbiological samples | 20036 | 4 bottles 100 ml | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED (EP) | For the dilution of microbiological samples | 20100 | 4 bottles 100 ml TP | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED (EP) | For the dilution of microbiological samples | 20008 | 4 bottles 90 ml | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED (EP) | For the dilution of microbiological samples | 20032 | 10 glass tubes | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED (EP) | For the dilution of microbiological samples | 20436 | 10 glass tubes 9 ml | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED (EP) | For the dilution of microbiological samples | 20032,5 | 50 glass tubes | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED (EP) | For the dilution of microbiological samples | 20436,5 | 50 glass tubes 9 ml | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED (EP) | For the dilution of microbiological samples | 20035 | 20 polystyrene tubes | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED (EP) | For the dilution of microbiological samples | 20163 | 4 bottles 200 ml | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED (EP) | For the dilution of microbiological samples | 20099 | 4 bottles 250 ml TP | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED (EP) | For the dilution of microbiological samples | 20990 | 4 bottles 300 ml | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED (EP) | For the dilution of microbiological samples | 20991 | 4 bottles 300 ml TP | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED (EP) | For the dilution of microbiological samples | 20760 | 4 bottles 500 ml | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED (EP) | For the dilution of microbiological samples | 20098 | 4 bottles 500 ml TP | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED (EP) | For the dilution of microbiological samples | 20435 | 4 bottles 1 lt | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED (EP) | For the dilution of microbiological samples | 20960 | 4 bottles 1 lt TP | 270 days | 8-25 | |
| PEPTONE WATER, BUFFERED (EP) | For the dilution of microbiological samples | 6194 | Dehydrated 500 gr | 3 years | 8-25 | |
| PEPTONE WATER, BUFFERED (EP) + LTH | For the dilution of microbiological samples and for the mirobiological test of pharmaceutical non sterile products | 20772 | 4 bottles 100 ml | 180 days | 4-8 | |
| PEPTONE WATER, BUFFERED (EP) + LTH | For the dilution of microbiological samples and for the mirobiological test of pharmaceutical non sterile products | 21058 | 4 bottles 90 ml | 180 days | 4-8 | |
| PEPTONE WATER, BUFFERED (EP) + LTH | For the dilution of microbiological samples and for the mirobiological test of pharmaceutical non sterile products | 21024 | 4 bottles 400 ml | 180 days | 4-8 | |
| PEPTONE WATER, BUFFERED (EP) + LTH | For the dilution of microbiological samples and for the mirobiological test of pharmaceutical non sterile products | 21187 | 4 bottles 1000 ml | 180 days | 4-8 | |
| POLIPEPTONE | For use in microbiology | 6413 | Dehydrated 500 gr | 3 years | 8-25 | |
| POTASSIUM PHOSPHATE BIBASIC 20 gr/l pH 7,5 | | 21367 | 2 bags 3 lt | 210 days | 8-25 | |
| POTASSIUM PHOSPHATE BIBASIC 20 gr/l pH 7,5 | | 21368 | 2 bags 5 lt | 210 days | 8-25 | |
| POTASSIUM PHOSPHATE MONOBASIC 20 gr/l pH 7,5 | | 21202 | 2 bags 3 lt | 210 days | 8-25 | |
| POTASSIUM PHOSPHATE MONOBASIC 20 gr/l pH 7,5 | | 21203 | 2 bags 5 lt | 210 days | 8-25 | |
| P.P.L.O. | For the culture of Mycoplasma spp. | 20097 | 20 plates 90 mm | 180 days | 4-8 | |
| P.P.L.O. AGAR BASE | For the culture of Mycoplasma spp. | 6616 | Dehydrated 500 gr | 3 years | 8-25 | |
| P.P.L.O. AGAR BASE – REQUIRED SUPPLEMENT | | 1497 | 1 x 100 ml / HORSE SERUM | 1 year | 4-8 | |
| PROTOTHECA ISOLATION MEDIUM (P.I.M.) | For the isolation of Prototheca spp. | 20946 | 20 plates 90 mm | 180 days | 4-8 | |
| PROTOTHECA ISOLATION MEDIUM (P.I.M.) AGAR BASE | For the isolation of Prototheca spp. | 6904 | Dehydrated 500 gr | 3 years | 8-25 | |
| PROTOTHECA ISOLATION MEDIUM (P.I.M.) AGAR BASE – REQUIRED SUPPLEMENT | | 6550 | 5 + 5 vials x 500 ml / PIM SUPPLEMENT | 3 years | 4-8 | |

| | | | | | | |
|--|--|-------|---|----------|------|--|
| PSEUDOMONAS CN ISO 16266 | For the identification of Pseudomonas aeruginosa by membrane filter method | 20065 | 20 plates 90 mm | 180 days | 4-8 | |
| PSEUDOMONAS CN ISO 16266 | For the identification of Pseudomonas aeruginosa by membrane filter method | 127 | 40 plates 60 mm | 180 days | 4-8 | |
| PSEUDOMONAS CN ISO 16266 | For the identification of Pseudomonas aeruginosa by membrane filter method | 20037 | 4 bottles 100 ml | 180 days | 4-8 | |
| PSEUDOMONAS CN AGAR BASE ISO 16266 | For the identification of Pseudomonas aeruginosa by membrane filter method | 6204 | Dehydrated 500 gr | 3 years | 8-25 | |
| PSEUDOMONAS CN AGAR BASE ISO 16266 – REQUIRED SUPPLEMENT | | 6316 | 1 x 1 lt / GLYCEROL | 3 years | 8-25 | |
| PSEUDOMONAS CFC ISO 13720 | For the selective isolation of Pseudomonas spp. | 20816 | 20 plates 90 mm | 180 days | 4-8 | |
| PSEUDOMONAS CFC ISO 13720 | For the selective isolation of Pseudomonas spp. | 20755 | 40 plates 60 mm | 180 days | 4-8 | |
| PSEUDOMONAS PP ISO / TS 11059 | For the count of Pseudomonas spp. | 20832 | 20 plates 90 mm | 180 days | 4-8 | |
| PSEUDOMONAS AGAR BASE | For the count of Pseudomonas spp. | 6854 | Dehydrated 500 gr | 3 years | 8-25 | |
| PSEUDOMONAS AGAR BASE – REQUIRED SUPPLEMENT | | 6537 | 10 vials x 500 ml / PSEUDOMONAS PP SUPPLEMENT ISO / TS 11059 E GLYCEROL | 3 years | 4-8 | |
| PSEUDOMONAS AGAR BASE – REQUIRED SUPPLEMENT | | 6520 | 10 vials x 500 ml / PSEUDOMONAS CFC SUPPLEMENT AND GLYCEROL ISO 13720 | 3 years | 4-8 | |
| PSEUDOMONAS SELECTIVE (EP) | For the identification of Pseudomonas aeruginosa by membrane filter method | 1016 | 20 plates 90 mm | 240 days | 8-25 | |
| PSEUDOMONAS SELECTIVE (EP) | For the identification of Pseudomonas aeruginosa by membrane filter method | 2216 | 40 plates 60 mm | 240 days | 8-25 | |
| PSEUDOMONAS SELECTIVE (EP) | For the identification of Pseudomonas aeruginosa by membrane filter method | 4016 | 40 contact 55 mm | 240 days | 8-25 | |
| PSEUDOMONAS SELECTIVE (EP) | For the identification of Pseudomonas aeruginosa by membrane filter method | 2516 | 40 contact s/m | 240 days | 8-25 | |
| PSEUDOMONAS SELECTIVE (EP) | For the identification of Pseudomonas aeruginosa by membrane filter method | 1216 | 4 bottles 100 ml | 240 days | 8-25 | |
| R2A (EP) | For the total aerobic bacterial count in drinking treated waters | 1045 | 20 plates 90 mm | 210 days | 8-25 | |
| R2A (EP) | For the total aerobic bacterial count in drinking treated waters | 2245 | 40 plates 60 mm | 210 days | 8-25 | |
| R2A (EP) | For the total aerobic bacterial count in drinking treated waters | 6080 | Dehydrated 500 gr | 3 years | 8-25 | |
| RHAMNOSE (L) BROTH ISO 11290 | For the confirmation of Listeria spp. | 20208 | 10 glass tubes | 180 days | 8-25 | |
| RAPPAPORT VASSILIADIS SOY BROTH (RVS) (EP) | For the enrichment of Salmonella spp. | 20308 | 10 glass tubes | 180 days | 8-25 | |
| RAPPAPORT VASSILIADIS SOY BROTH (RVS) (EP) | For the enrichment of Salmonella spp. | 20307 | 4 bottles 100 ml | 180 days | 8-25 | |
| RAPPAPORT VASSILIADIS SOY BROTH (RVS) (EP) | For the enrichment of Salmonella spp. | 6180 | Dehydrated 500 gr | 3 years | 8-25 | |
| RAPPAPORT VASSILIADIS SOY BROTH ISO 6579 | For the enrichment of Salmonella spp. | 20416 | 10 glass tubes | 180 days | 8-25 | |
| RAPPAPORT VASSILIADIS SOY BROTH ISO 6579 | For the enrichment of Salmonella spp. | 20566 | 4 bottles 225 ml | 180 days | 8-25 | |
| RAPPAPORT VASSILIADIS SOY BROTH ISO 6579 | For the enrichment of Salmonella spp. | 20443 | 4 bottles 100 ml | 180 days | 8-25 | |
| RAPPAPORT VASSILIADIS SOY BROTH ISO 6579 | For the enrichment of Salmonella spp. | 20660 | 4 bottles 90 ml | 180 days | 8-25 | |
| RAPPAPORT VASSILIADIS SOY BROTH ISO 6579 | For the enrichment of Salmonella spp. | 6617 | Dehydrated 500 gr | 3 years | 8-25 | |
| RAPPAPORT MODIFIED SEMISOLID (MRSV) ISO 6579 | For the detection of mobile Salmonella | 20390 | 4 bottles 100 ml | 90 days | 4-8 | |
| RAPPAPORT MODIFIED SEMISOLID (MRSV) ISO 6579 | For the detection of mobile Salmonella | 20746 | 10 glass tubes | 90 days | 4-8 | |

| | | | | | | |
|--|---|--------|---|----------|------|--|
| RAPPAPORT MODIFIED SEMISOLID (MRSV) ISO 6579 | For the detection of mobile Salmonella | 6744 | Dehydrated 500 gr | 3 years | 8-25 | |
| RAPPAPORT MODIFIED SEMISOLID (MRSV) ISO 6579 – REQUIRED SUPPLEMENT | | 6538 | 10 vials x 500 ml / NOVOBIOCIN SUPPLEMENT | 3 years | 4-8 | |
| REINFORCED CLOSTRIDIAL AGAR | For the culture of Clostridium spp. and other anaerobic microorganisms | 20166 | 4 bottles 100 ml | 180 days | 4-8 | |
| REINFORCED CLOSTRIDIAL AGAR | For the culture of Clostridium spp. and other anaerobic microorganisms | 20325 | 20 plates 90 mm | 180 days | 4-8 | |
| REINFORCED CLOSTRIDIAL AGAR | For the culture of Clostridium spp. and other anaerobic microorganisms | 6619 | Dehydrated 500 gr | 3 years | 8-25 | |
| REINFORCED CLOSTRIDIAL MEDIUM (EP) | For the culture of Clostridium spp. and other anaerobic microorganisms | 124 | 4 bottles 100 ml | 180 days | 4-8 | |
| REINFORCED CLOSTRIDIAL MEDIUM (EP) | For the culture of Clostridium spp. and other anaerobic microorganisms | 6620 | Dehydrated 500 gr | 3 years | 8-25 | |
| ROGOSA | For the isolation and the enumeration of Lactobacilli | 1034 | 20 plates 90 mm | 180 days | 4-8 | |
| ROGOSA SL AGAR BASE | For the isolation and the enumeration of Lactobacilli | 6179 | Dehydrated 500 gr | 3 years | 8-25 | |
| ROSE BENGAL | For the isolation of yeast and fungi | 1046 A | 20 plates 90 mm | 180 days | 4-8 | |
| ROSE BENGAL | For the isolation of yeast and fungi | 4046 A | 40 contact 55 mm | 180 days | 4-8 | |
| ROSE BENGAL | For the isolation of yeast and fungi | 6209 | Dehydrated 500 gr | 3 years | 8-25 | |
| ROSE BENGAL + DICHLORAN ISO 21527 | For the count of yeasts and fungi in the food and feed with aw more than 0,95 | 20616 | 20 plates 90 mm | 180 days | 4-8 | |
| ROSE BENGAL + DICHLORAN ISO 21527 | For the count of yeasts and fungi in the food and feed with aw more than 0,95 | 20678 | 40 contact s/m | 180 days | 4-8 | |
| ROSE BENGAL + DICHLORAN ISO 21527 | For the count of yeasts and fungi in the food and feed with aw more than 0,95 | 20378 | 4 bottles 100 ml | 180 days | 4-8 | |
| ROSE BENGAL + DICHLORAN ISO 21527 | For the count of yeasts and fungi in the food and feed with aw more than 0,95 | 6623 | Dehydrated 500 gr | 3 years | 8-25 | |
| SABOURAUD DEXTROSE AGAR (EP) ISO 18416 | For the culture of yeasts and fungi | 1017 | 20 plates 90 mm | 210 days | 8-25 | |
| SABOURAUD DEXTROSE AGAR (EP) ISO 18416 | For the culture of yeasts and fungi | 2217 | 40 plates 60 mm | 210 days | 8-25 | |
| SABOURAUD DEXTROSE AGAR (EP) ISO 18416 | For the culture of yeasts and fungi | 4017 | 40 contact 55 mm | 210 days | 8-2 | |
| SABOURAUD DEXTROSE AGAR (EP) ISO 18416 | For the culture of yeasts and fungi | 2517 | 40 contact s/m | 210 days | 8-25 | |
| SABOURAUD DEXTROSE AGAR (EP) ISO 18416 | For the culture of yeasts and fungi | 1217 | 4 bottles 100 ml | 210 days | 8-25 | |
| SABOURAUD DEXTROSE AGAR (EP) ISO 18416 | For the culture of yeasts and fungi | 1117 | 10 glass tubes slant | 210 days | 8-25 | |
| SABOURAUD DEXTROSE AGAR (EP) ISO 18416 | For the culture of yeasts and fungi | 20639 | 4 bottles 500 ml | 210 days | 8-25 | |
| SABOURAUD DEXTROSE AGAR (EP) ISO 18416 | For the culture of yeasts and fungi | 20689 | 20 contact 90 mm s/m | 210 days | 8-25 | |
| SABOURAUD DEXTROSE AGAR (EP) ISO 18416 | For the culture of yeasts and fungi | 20770 | 4 bottles 200 ml | 210 days | 8-25 | |
| SABOURAUD DEXTROSE AGAR (EP) ISO 18416 | For the culture of yeasts and fungi | 20640 | 4 bottles 1000 ml | 210 days | 8-25 | |
| SABOURAUD DEXTROSE AGAR (EP) ISO 18416 | For the culture of yeasts and fungi | 6081 | Dehydrated 500 gr | 3 years | 8-25 | |
| SABOURAUD DEXTROSE UNICHIM | For the culture of yeasts and fungi | 20385 | 40 contact 55 mm | 210 days | 8-25 | |
| SABOURAUD DEXTROSE UNICHIM | For the culture of yeasts and fungi | 20373 | 40 contact s/m | 210 days | 8-25 | |
| SAB + CAF 50 ISO 18416 | For the isolation of yeast and fungi | 1018 A | 20 plates 90 mm | 180 days | 4-8 | |
| SAB + CAF 50 ISO 18416 | For the isolation of yeast and fungi | 2218 A | 40 plates 60 mm | 180 days | 4-8 | |
| SAB + CAF 50 ISO 18416 | For the isolation of yeast and fungi | 4018 A | 40 contact 55 mm | 180 days | 4-8 | |
| SAB + CAF 50 ISO 18416 | For the isolation of yeast and fungi | 2518 A | 40 contact s/m | 180 days | 4-8 | |

| | | | | | | |
|---------------------------------------|---|--------|----------------------|----------|------|--|
| SAB + CAF 50 ISO 18416 | For the isolation of yeast and fungi | 1218 A | 4 bottles 100 ml | 180 days | 4-8 | |
| SAB + CAF 50 ISO 18416 | For the isolation of yeast and fungi | 6210 | Dehydrated 500 gr | 3 years | 8-25 | |
| SAB + CAF 500 | For the isolation of yeast and fungi | 1018 | 20 plates 90 mm | 180 days | 4-8 | |
| SAB + CAF 500 | For the isolation of yeast and fungi | 2218 | 40 plates 60 mm | 180 days | 4-8 | |
| SAB + CAF 500 | For the isolation of yeast and fungi | 1218 | 4 bottles 100 ml | 180 days | 4-8 | |
| SAB + CAF 500 | For the isolation of yeast and fungi | 20826 | 10 glass tubes 15 ml | 180 days | 4-8 | |
| SAB + CAF 500 | For the isolation of yeast and fungi | 6082 | Dehydrated 500 gr | 3 years | 8-25 | |
| SAB + CAF 50 + GENTAMICIN | For the isolation of yeast and fungi | 1086 | 20 plates 90 mm | 180 days | 4-8 | |
| SABOURAUD + LECITHIN + Tween 80 | For the culture of yeasts and fungi | 4018 | 40 contact 55 mm | 180 days | 4-8 | |
| SABOURAUD + LECITHIN + Tween 80 | For the culture of yeasts and fungi | 20153 | 4 bottles 100 ml | 180 days | 4-8 | |
| SABOURAUD + CAF 50 + CEX | For the isolation of pathogenic fungi | 1018 C | 20 plates 90 mm | 180 days | 4-8 | |
| SABOURAUD + CAF 50 + CEX | For the isolation of pathogenic fungi | 1180 C | 10 glass tubes | 180 days | 4-8 | |
| SABOURAUD + CAF 50 + CEX | For the isolation of pathogenic fungi | 6083 | Dehydrated 500 gr | 3 years | 4-8 | |
| SABOURAUD + CEX | For the isolation of pathogenic fungi | 20186 | 20 plates 90 mm | 180 days | 4-8 | |
| SABOURAUD + CEX | For the isolation of pathogenic fungi | 6258 | Dehydrated 500 gr | 3 years | 4-8 | |
| SAB + LECITHIN + Tween 80 + CAF 50 | For the isolation of yeast and fungi | 4018 B | 40 contact 55 mm | 180 days | 4-8 | |
| SABOURAUD BROTH (EP) | For the culture of yeasts and fungi | 1163 | 10 glass tubes | 210 days | 8-25 | |
| SABOURAUD BROTH (EP) | For the culture of yeasts and fungi | 1263 | 4 bottles 100 ml | 210 days | 8-25 | |
| SABOURAUD BROTH (EP) | For the culture of yeasts and fungi | 1863 | 20 polystyrene tubes | 210 days | 8-25 | |
| SABOURAUD BROTH (EP) | For the culture of yeasts and fungi | 20641 | 4 bottles 500 ml | 210 days | 8-25 | |
| SABOURAUD BROTH (EP) | For the culture of yeasts and fungi | 20642 | 4 bottles 1000 ml | 210 days | 8-25 | |
| SABOURAUD BROTH (EP) | For the culture of yeasts and fungi | 6084 | Dehydrated 500 gr | 3 years | 8-25 | |
| SABOURAUD + CAF 50 BROTH | For the isolation of yeast and fungi | 1263 A | 4 bottles 100 ml | 180 days | 4-8 | |
| SABOURAUD + CAF 50 + GENTAMICIN BROTH | For the isolation of yeast and fungi | 1186 | 10 glass tubes | 180 days | 4-8 | |
| SABOURAUD FLUID (USP) | For the culture of yeasts and fungi | 1131 | 10 glass tubes | 210 days | 8-25 | |
| SABOURAUD FLUID (USP) | For the culture of yeasts and fungi | 1831 | 20 polystyrene tubes | 210 days | 8-25 | |
| SABOURAUD FLUID (USP) | For the culture of yeasts and fungi | 6085 | Dehydrated 500 gr | 3 years | 8-25 | |
| SABOURAUD FLUID + Tween 80 | For the culture of yeasts and fungi | 1131 A | 10 glass tubes | 210 days | 8-25 | |
| SOLUTION 0,90% NaCl | For the dilution of microbiological samples | 20291 | 10 glass tubes | 270 days | 8-25 | |
| SOLUTION 0,90% NaCl | For the dilution of microbiological samples | 20411 | 10 glass tubes 5 ml | 270 days | 8-25 | |
| SOLUTION 0,90% NaCl | For the dilution of microbiological samples | 20158 | 10 glass tubes 9 ml | 270 days | 8-25 | |
| SOLUTION 0,90% NaCl | For the dilution of microbiological samples | 20159 | 4 bottles 90 ml | 270 days | 8-25 | |
| SOLUTION 0,90% NaCl | For the dilution of microbiological samples | 20794 | 4 bottles 100 ml | 270 days | 8-25 | |
| SOLUTION 0,90% NaCl | For the dilution of microbiological samples | 20936 | 2 bags 3 lt | 270 days | 8-25 | |
| SOLUTION 0,85% NaCl ISO 6579 | For the maintenance of the bacterial strains | 1185 | 10 glass tubes | 270 days | 8-25 | |
| SOLUTION 0,85% NaCl ISO 6579 | For the maintenance of the bacterial strains | 1285 | 4 bottles 100 ml | 270 days | 8-25 | |
| SCHAEDLER BLOOD | For the culture of anaerobic bacteria | 1059 | 20 plates 90 mm | 90 days | 4-8 | |
| SCHAEDLER SELECTIVE BLOOD | For the culture of Gram-negative anaerobic bacteria | 1060 | 20 plates 90 mm | 90 days | 4-8 | |
| SCHAEDLER SELECTIVE BLOOD CNA | For the culture of Gram-positive anaerobic bacteria | 1060 A | 20 plates 90 mm | 90 days | 4-8 | |

| | | | | | | |
|-------------------------|--|--------|----------------------|----------|------|--|
| SCHAEDLER AGAR BASE | For the culture of anaerobic bacteria | 6088 | Dehydrated 500 gr | 3 years | 8-25 | |
| SELENITE BROTH | For the selective enrichment of <i>Salmonella</i> spp. | 1164 | 10 glass tubes | 210 days | 4-8 | |
| SELENITE BROTH | For the selective enrichment of <i>Salmonella</i> spp. | 1864 | 20 polystyrene tubes | 210 days | 4-8 | |
| SELENITE BROTH | For the selective enrichment of <i>Salmonella</i> spp. | 20165 | 10 glass tubes 20 ml | 210 days | 4-8 | |
| SELENITE BROTH | For the selective enrichment of <i>Salmonella</i> spp. | 1864 | 20 polystyrene tubes | 210 days | 4-8 | |
| SELENITE BROTH | For the selective enrichment of <i>Salmonella</i> spp. | 20165 | 10 glass tubes 20 ml | 210 days | 4-8 | |
| SELENITE BROTH | For the selective enrichment of <i>Salmonella</i> spp. | 1164 C | 10 glass tubes 5 ml | 210 days | 4-8 | |
| SELENITE BROTH | For the selective enrichment of <i>Salmonella</i> spp. | 1264 | 4 bottles 100 ml | 210 days | 4-8 | |
| SELENITE BROTH | For the selective enrichment of <i>Salmonella</i> spp. | 20565 | 4 bottles 225 ml | 210 days | 4-8 | |
| SELENITE BROTH | For the selective enrichment of <i>Salmonella</i> spp. | 6091 | Dehydrated 500 gr | 3 years | 4-8 | |
| SELENITE CYSTINE BROTH | For the selective enrichment of <i>Salmonella</i> spp. and some strains of <i>Shigella</i> | 1164 A | 10 glass tubes | 210 days | 4-8 | |
| SELENITE CYSTINE BROTH | For the selective enrichment of <i>Salmonella</i> spp. and some strains of <i>Shigella</i> | 1164 B | 10 glass tubes 20 ml | 210 days | 4-8 | |
| SELENITE CYSTINE BROTH | For the selective enrichment of <i>Salmonella</i> spp. and some strains of <i>Shigella</i> | 1864 A | 20 polystyrene tubes | 210 days | 4-8 | |
| SELENITE CYSTINE BROTH | For the selective enrichment of <i>Salmonella</i> spp. and some strains of <i>Shigella</i> | 1264 A | 4 bottles 100 ml | 210 days | 4-8 | |
| SELENITE CYSTINE BROTH | For the selective enrichment of <i>Salmonella</i> spp. and some strains of <i>Shigella</i> | 1264 A | 40 bottles 100 ml | 210 days | 4-8 | |
| SELENITE CYSTINE BROTH | For the selective enrichment of <i>Salmonella</i> spp. and some strains of <i>Shigella</i> | 6097 | Dehydrated 500 gr | 3 years | 4-8 | |
| SERUM TELLURITE | For the isolation of <i>Corynebacterium</i> spp. | 1085 | 20 plates 90 mm | 180 days | 4-8 | |
| SF BROTH | For the culture of fecal Streptococci | 1165 | 10 glass tubes | 270 days | 8-25 | |
| SF BROTH | For the culture of fecal Streptococci | 1865 | 20 polystyrene tubes | 270 days | 8-25 | |
| SIMMONS CITRATE AGAR | For the determination of Enterobacteria according to the use of citrate | 6725 | Dehydrated 500 gr | 3 years | 8-25 | |
| SKIM MILK AGAR ISO 6610 | For total bacteria count in the milk and dairy products | 20180 | 20 plates 90 mm | 240 days | 4-8 | |
| SKIM MILK AGAR ISO 6610 | For total bacteria count in the milk and dairy products | 20197 | 10 glass tubes 15 ml | 240 days | 4-8 | |
| SKIM MILK AGAR ISO 6610 | For total bacteria count in the milk and dairy products | 6626 | Dehydrated 500 gr | 3 years | 8-25 | |
| SMSA | For the isolation of <i>Pseudomonas</i> spp. | 3006 | 20 plates 90 mm | 180 days | 4-8 | |
| SOB BROTH | For the culture of recombinant strains of <i>Escherichia coli</i> | 6700 | Dehydrated 500 gr | 3 years | 8-25 | |
| SOC BROTH | For the culture of recombinant strains of <i>Escherichia coli</i> | 20096 | 4 bottles 100 ml | 270 days | 4-8 | |
| SODIUM ACETATE AGAR | For the determination of Enterobacteria according to the use of acetate | 6727 | Dehydrated 500 gr | 3 years | 8-25 | |
| SOY LECITHIN | | 6325 | Dehydrated 1000 gr | 3 years | 4-8 | |
| SPS | For the isolation of <i>Clostridium perfringens</i> | 1044 | 20 plates 90 mm | 150 days | 4-8 | |
| SPS | For the isolation of <i>Clostridium perfringens</i> | 2244 | 40 plates 60 mm | 150 days | 4-8 | |
| SPS | For the isolation of <i>Clostridium perfringens</i> | 20375 | 10 glass tubes 15 ml | 150 days | 4-8 | |
| SPS | For the isolation of <i>Clostridium perfringens</i> | 1144 | 10 glass tubes 22 ml | 150 days | 4-8 | |
| SPS | For the isolation of <i>Clostridium perfringens</i> | 1244 | 4 bottles 100 ml | 150 days | 4-8 | |
| SPS | For the isolation of <i>Clostridium perfringens</i> | 6094 | Dehydrated 500 gr | 3 years | 4-8 | |

| | | | | | | |
|---|---|---------|------------------------------------|----------|------|--|
| SSDC ISO 10273 | For the detection of Yersinia enterocolitica | 20810 | 20 plates 90 mm | 210 days | 8-25 | |
| SSDC ISO 10273 | For the detection of Yersinia enterocolitica | 6852 | Dehydrated 500 gr | 3 years | 8-25 | |
| SS SALMONELLA - SHIGELLA AGAR | For the isolation of Salmonella and Shigella | 1020 | 20 plates 90 mm | 210 days | 8-25 | |
| SS SALMONELLA - SHIGELLA AGAR | For the isolation of Salmonella and Shigella | 1220 | 4 bottles 100 ml | 210 days | 8-25 | |
| SS SALMONELLA - SHIGELLA AGAR | For the isolation of Salmonella and Shigella | 6087 | Dehydrated 500 gr | 3 years | 8-25 | |
| STERILE DEIONIZED WATER | For use in microbiology | 20784 | 10 glass tubes 5 ml | 365 days | 8-25 | |
| STERILE DEIONIZED WATER | For use in microbiology | 20278 | 4 bottles 100 ml | 365 days | 8-25 | |
| STERILE DEIONIZED WATER | For use in microbiology | 20888 | 4 bottles 400 ml | 365 days | 8-25 | |
| STERILE DEIONIZED WATER | For use in microbiology | 20930 | 4 bottles 1000 ml | 365 days | 8-25 | |
| STERILE DEIONIZED WATER | For use in microbiology | 20941 | 4 bottles 1000 ml TP | 365 days | 8-25 | |
| STERILE DEIONIZED WATER | For use in microbiology | 20942 | 4 bottles 500 ml TP | 365 days | 8-25 | |
| STERILE DEIONIZED WATER | For use in microbiology | 20934 | 2 bags 3 lt | 365 days | 8-25 | |
| SUCROSE | | 6437 | 500 gr | 3 years | 8-25 | |
| TAYLORELLA EQUIGENITALIS AGAR BASE | For the culture of Taylorella equigenitalis | 6732 | Dehydrated 500 gr | 3 years | 8-25 | |
| TCBS | For the isolation of Vibrio spp. | 1043 | 20 plates 90 mm | 180 days | 4-8 | |
| TCBS | For the isolation of Vibrio spp. | 6098 | Dehydrated 500 gr | 3 years | 8-25 | |
| TERGITOL 7 | For the count of Coliforms in the waters by filtering membrane method | 2249 | 40 plates 60 mm | 180 days | 4-8 | |
| TCBS | For the isolation of Vibrio spp. | 4049 | 40 contact 55 mm | 180 days | 4-8 | |
| TERGITOL 7 + TTC (ISO 9308) | For the count of Coliforms in the waters by filtering membrane method | 2249 A | 40 plates 60 mm | 180 days | 4-8 | |
| TERGITOL 7 + TTC (ISO 9308) | For the count of Coliforms in the waters by filtering membrane method | 4049 A | 40 contact 55 mm | 180 days | 4-8 | |
| TERGITOL 7 + TTC (ISO 9308) | For the count of Coliforms in the waters by filtering membrane method | 20142 | 20 plates 90 mm | 180 days | 4-8 | |
| TERGITOL 7 AGAR BASE (ISO 9308) | For the count of Coliforms in the waters by filtering membrane method | 6218 | Dehydrated 500 gr | 3 years | 8-25 | |
| TERGITOL 7 AGAR BASE (ISO 9308) – REQUIRED SUPPLEMENT | | 6509 | 10 vials x 500 ml / TTC SUPPLEMENT | 3 years | 4-8 | |
| THAYER MARTIN MODIFIED | For the isolation of Neisseria spp. | 1031 | 20 plates 90 mm | 180 days | 4-8 | |
| THIOGLYCOLLATE BROTH (NIH-USP) | For sterility test | 6216 | Dehydrated 500 gr | 3 years | 8-25 | |
| THIOGLYCOLLATE FLUID USP | For the culture of aerobic and anaerobic bacteria | 1167 | 10 glass tubes | 180 days | 8-25 | |
| THIOGLYCOLLATE FLUID USP | For the culture of aerobic and anaerobic bacteria | 20066 | 10 glass tubes 15 ml | 180 days | 8-25 | |
| THIOGLYCOLLATE FLUID USP | For the culture of aerobic and anaerobic bacteria | 1167,5 | 50 glass tubes | 180 days | 8-25 | |
| THIOGLYCOLLATE FLUID USP | For the culture of aerobic and anaerobic bacteria | 20066,5 | 50 glass tubes 15 ml | 180 days | 8-25 | |
| THIOGLYCOLLATE FLUID USP | For the culture of aerobic and anaerobic bacteria | 1867 | 20 polystyrene tubes | 180 days | 8-25 | |
| THIOGLYCOLLATE FLUID USP | For the culture of aerobic and anaerobic bacteria | 1267 | 4 bottles 100 ml | 180 days | 8-25 | |
| THIOGLYCOLLATE FLUID USP | For the culture of aerobic and anaerobic bacteria | 126 | 4 bottles 100 ml TP | 180 days | 8-25 | |
| THIOGLYCOLLATE FLUID USP | For the culture of aerobic and anaerobic bacteria | 6100 | Dehydrated 500 gr | 3 years | 8-25 | |
| TODD HEWITT BROTH | For the culture of Streptococci | 1168 | 10 glass tubes | 270 days | 8-25 | |
| TODD HEWITT BROTH | For the culture of Streptococci | 1868 | 20 polystyrene tubes | 270 days | 8-25 | |
| TODD HEWITT BROTH | For the culture of Streptococci | 6102 | Dehydrated 500 gr | 3 years | 8-25 | |
| TRICHOMONAS SELECTIVE | For the isolation of Trichomonas vaginalis | 1173 | 10 glass tubes 5 ml | 150 days | 4-8 | |

| | | | | | | |
|---------------------------------------|--|-------|----------------------------|----------|------|--|
| TRICHOMONAS SELECTIVE | For the isolation of Trichomonas vaginalis | 1873 | 20 polystyrene tubes 5 ml | 150 days | 4-8 | |
| TRIPLE SUGAR IRON AGAR (TSI) (EP) | For the identification and differentiation of Enterobacteria | 1138 | 10 glass tubes slant | 210 days | 4-8 | |
| TRIPLE SUGAR IRON AGAR (TSI) (EP) | For the identification and differentiation of Enterobacteria | 1838 | 20 polystyrene tubes slant | 210 days | 4-8 | |
| TRIPLE SUGAR IRON AGAR (TSI) (EP) | For the identification and differentiation of Enterobacteria | 6103 | Dehydrated 500 gr | 3 years | 8-25 | |
| TRIPLE SUGAR IRON AGAR (TSI) ISO 6579 | For the identification and differentiation of Enterobacteria | 20287 | 20 plates 90 mm | 210 days | 8-25 | |
| TRIPLE SUGAR IRON AGAR (TSI) ISO 6579 | For the identification and differentiation of Enterobacteria | 20724 | 10 glass tubes slant | 210 days | 8-25 | |
| TRIPLE SUGAR IRON AGAR (TSI) ISO 6579 | For the identification and differentiation of Enterobacteria | 6276 | Dehydrated 500 gr | 3 years | 8-25 | |
| TRYPTOPHAN BROTH (ISO 9308 E 6579) | For the determination of E. Coli and other Coliforms according to indole production | 20140 | 20 polystyrene tubes 3 ml | 180 days | 8-25 | |
| TRYPTOPHAN BROTH (ISO 9308 E 6579) | For the determination of E. Coli and other Coliforms according to indole production | 20015 | 4 bottles 100 ml | 180 days | 8-25 | |
| TRYPTOPHAN BROTH (ISO 9308 E 6579) | For the determination of E. Coli and other Coliforms according to indole production | 20143 | 10 glass tubes 3 ml | 180 days | 8-25 | |
| TRYPTOPHAN BROTH (ISO 9308 E 6579) | For the determination of E. Coli and other Coliforms according to indole production | 20016 | 10 glass tubes 5 ml | 180 days | 8-25 | |
| TRYPTOPHAN BROTH (ISO 9308 E 6579) | For the determination of E. Coli and other Coliforms according to indole production | 6633 | Dehydrated 500 gr | 3 years | 8-25 | |
| TRYPTONE | For use in microbiology | 6416 | Dehydrated 500 gr | 3 years | 8-25 | |
| TRYPTONE BILE AGAR (ISO 9308) | For the determination and the enumeration of E. Coli and other Coliforms in water and food | 20146 | 20 plates 90 mm | 270 days | 8-25 | |
| TRYPTONE BILE AGAR (ISO 9308) | For the determination and the enumeration of E. Coli and other Coliforms in water and food | 20490 | 40 plates 60 mm | 270 days | 8-25 | |
| TRYPTONE BILE AGAR (ISO 9308) | For the determination and the enumeration of E. Coli and other Coliforms in water and food | 20561 | 4 bottles 100 ml | 270 days | 8-25 | |
| TRYPTONE BILE AGAR (ISO 9308) | For the determination and the enumeration of E. Coli and other Coliforms in water and food | 6294 | Dehydrated 500 gr | 3 years | 8-25 | |
| TRYPTONE SOY AGAR ISO 9308 | For the enumeration of Escherichia Coli and other Coliforms | 20420 | 20 plates 90 mm | 240 days | 8-25 | |
| TRYPTONE SOY AGAR ISO 9308 | For the enumeration of Escherichia Coli and other Coliforms | 20419 | 40 plates 60 mm | 240 days | 8-25 | |
| TRYPTONE SOY AGAR ISO 9308 | For the enumeration of Escherichia Coli and other Coliforms | 6277 | Dehydrated 500 gr | 3 years | 8-25 | |
| TRYPTOSE | For use in microbiology | 6417 | Dehydrated 500 gr | 3 years | 8-25 | |
| TRYPTOSE AGAR | For the culture of fastidious microorganisms especially Brucella spp. | 1026 | 20 plates 90 mm | 270 days | 8-25 | |
| TRYPTOSE AGAR | For the culture of fastidious microorganisms especially Brucella spp. | 1126 | 10 glass tubes slant | 270 days | 8-25 | |
| TRYPTOSE AGAR | For the culture of fastidious microorganisms especially Brucella spp. | 1826 | 20 polystyrene tubes slant | 270 days | 8-25 | |

| | | | | | | |
|---|--|--------|----------------------------|----------|------|--|
| TRYPTOSE AGAR | For the culture of fastidious microorganisms especially <i>Brucella</i> spp. | 1226 | 4 bottles 100 ml | 270 days | 8-25 | |
| TRYPTOSE AGAR | For the culture of fastidious microorganisms especially <i>Brucella</i> spp. | 6631 | Dehydrated 500 gr | 3 years | 8-25 | |
| TRYPTOSE BROTH | For the culture of fastidious microorganisms especially <i>Brucella</i> spp. | 1170 | 10 glass tubes | 270 days | 8-25 | |
| TRYPTOSE BROTH | For the culture of fastidious microorganisms especially <i>Brucella</i> spp. | 1870 | 20 polystyrene tubes | 270 days | 8-25 | |
| TRYPTOSE BROTH | For the culture of fastidious microorganisms especially <i>Brucella</i> spp. | 1270 | 4 bottles 100 ml | 270 days | 8-25 | |
| TRYPTOSE BROTH | For the culture of fastidious microorganisms especially <i>Brucella</i> spp. | 6632 | Dehydrated 500 gr | 3 years | 8-25 | |
| TSA (EP) | For the culture of fastidious bacteria and not | 1025 | 20 plates 90 mm | 240 days | 8-25 | |
| TSA (EP) | For the culture of fastidious bacteria and not | 2225 | 40 plates 60 mm | 240 days | 8-25 | |
| TSA (EP) | For the culture of fastidious bacteria and not | 4025 | 40 contact 55 mm | 240 days | 8-25 | |
| TSA (EP) | For the culture of fastidious bacteria and not | 2525 | 40 contact s/m | 240 days | 8-25 | |
| TSA (EP) | For the culture of fastidious bacteria and not | 20244 | 20 contact 90 mm | 240 days | 8-25 | |
| TSA (EP) | For the culture of fastidious bacteria and not | 1325 | plates10 plates 150 mm | 240 days | 8-25 | |
| TSA (EP) | For the culture of fastidious bacteria and not | 20425 | 10 glass tubes 15 ml | 240 days | 8-25 | |
| TSA (EP) | For the culture of fastidious bacteria and not | 1125 | 10 glass tubes slant | 240 days | 8-25 | |
| TSA (EP) | For the culture of fastidious bacteria and not | 1825 | 20 polystyrene tubes slant | 240 days | 8-25 | |
| TSA (EP) | For the culture of fastidious bacteria and not | 1225 | 4 bottles 100 ml | 240 days | 8-25 | |
| TSA (EP) | For the culture of fastidious bacteria and not | 20526 | 4 bottles 500 ml | 240 days | 8-25 | |
| TSA (EP) | For the culture of fastidious bacteria and not | 20527 | 4 bottles 250 ml | 240 days | 8-25 | |
| TSA (EP) | For the culture of fastidious bacteria and not | 20771 | 4 bottles 200 ml | 240 days | 8-25 | |
| TSA (EP) | For the culture of fastidious bacteria and not | 6105 | Dehydrated 500 gr | 3 years | 8-25 | |
| TSA + LECITHIN + TWEEN 80 | For the culture of fastidious bacteria and not | 20154 | 4 bottles 100 ml | 210 days | 4-8 | |
| TSA + LECITHIN + TWEEN 80 | For the culture of fastidious bacteria and not | 4025 A | 40 contact 55 mm | 210 days | 4-8 | |
| TSA + LECITHIN + TWEEN 80 | For the culture of fastidious bacteria and not | 2525 A | 40 contact s/m | 210 days | 4-8 | |
| TSA + LECITHIN + TWEEN 80 | For the culture of fastidious bacteria and not | 20155 | 4 bottles 400 ml | 210 days | 4-8 | |
| TSA + LECITHIN + TWEEN 80 | For the culture of fastidious bacteria and not | 20245 | 20 contact 90 mm | 210 days | 4-8 | |
| TSA + LECITHIN + TWEEN 80 | For the culture of fastidious bacteria and not | 20242 | 20 plates 90 mm | 210 days | 4-8 | |
| TSA + LECITHIN + TWEEN 80 + PENASI 20000 u.i./l | For the culture of fastidious bacteria and not | 20275 | 40 contact 55 mm | 90 days | 4-8 | |
| TSA + LECITHIN + TWEEN 80 + SODIUM THIOSULPHATE + HISTIDINE | For the culture of fastidious bacteria and not | 4025 B | 40 contact 55 mm | 210 days | 4-8 | |
| TSA + PENASI 20000 u.i./l | For the culture of fastidious bacteria and not | 20262 | 20 plates 90 mm | 90 days | 4-8 | |
| TSA + SHEEP BLOOD | General use media and study of haemolytic reactions | 1021 | 20 plates 90 mm | 90 days | 4-8 | |
| TSA + SHEEP BLOOD | General use media and study of haemolytic reactions | 1321 | plates10 plates 150 mm | 90 days | 4-8 | |
| TSA + SHEEP BLOOD 5% (EP, USDA, FSIS MIG 8.07) | For the culture of fastidious bacteria and not | 20230 | 20 plates 90 mm | 90 days | 4-8 | |
| TSA + TTC | For the culture of fastidious bacteria and not | 21362 | 40 contact 55 mm | 180 days | 4-8 | |
| TSB (EP) | For the culture of fastidious bacteria and not | 1269 | 4 bottles 100 ml | 240 days | 8-25 | |

| | | | | | | |
|---|---|--|---|----------|------|--|
| TSB (EP) | For the culture of fastidious bacteria and not | 1269,4 | 40 bottles 100 ml | 240 days | 8-25 | |
| TSB (EP) | For the culture of fastidious bacteria and not | 125 | 4 bottles 100 ml TP | 240 days | 8-25 | |
| TSB (EP) | For the culture of fastidious bacteria and not | 1169 | 10 glass tubes | 240 days | 8-25 | |
| TSB (EP) | For the culture of fastidious bacteria and not | 1169 A | 10 glass tubes 15 ml | 240 days | 8-25 | |
| TSB (EP) | For the culture of fastidious bacteria and not | 1169,5 | 50 glass tubes | 240 days | 8-25 | |
| TSB (EP) | For the culture of fastidious bacteria and not | 1169 A.50 | 50 glass tubes 15 ml | 240 days | 8-25 | |
| TSB (EP) | For the culture of fastidious bacteria and not | 20646 | 4 bottles 1000 ml | 240 days | 8-25 | |
| TSB (EP) | For the culture of fastidious bacteria and not | 20948 | 4 bottles 1000 ml TP | 240 days | 8-25 | |
| TSB (EP) | For the culture of fastidious bacteria and not | 20815 | 4 bottles 90 ml | 240 days | 8-25 | |
| TSB (EP) | For the culture of fastidious bacteria and not | 20815,4 | 40 bottles 90 ml | 240 days | 8-25 | |
| TSB (EP) | For the culture of fastidious bacteria and not | 20645 | 4 bottles 500 ml | 240 days | 8-25 | |
| TSB (EP) | For the culture of fastidious bacteria and not | 20950 | 4 bottles 500 ml TP | 240 days | 8-25 | |
| TSB (EP) | For the culture of fastidious bacteria and not | 6217 | Dehydrated 500 gr | 3 years | 8-25 | |
| TSB MODIFIED (ISO 16654) | For the selective enrichment of E. Coli O157:h7 | 20549 | 10 glass tubes | 90 days | 4-8 | |
| TSB MODIFIED (ISO 16654) | For the selective enrichment of E. Coli O157:h7 | 20723 | 10 glass tubes 9 ml | 90 days | 4-8 | |
| TSB MODIFIED (ISO 16654) | For the selective enrichment of E. Coli O157:h7 | 20875 | 4 bottles 90 ml | 90 days | 4-8 | |
| TSB MODIFIED (ISO 16654) | For the selective enrichment of E. Coli O157:h7 | 20479 | 4 bottles 225 ml | 90 days | 4-8 | |
| TSB MODIFIED (ISO 16654) | For the selective enrichment of E. Coli O157:h7 | 20550 | 4 bottles 100 ml | 90 days | 4-8 | |
| TSB MODIFIED BROTH BASE (ISO 16654) | For the selective enrichment of E. Coli O157:h7 | 6651 | Dehydrated 500 gr | 3 years | 8-25 | |
| TSB MODIFIED BROTH BASE (ISO 16654) – REQUIRED SUPPLEMENT | | 6526 | 10 vials x 500 ml / NOVOBIOCIN SUPPLEMENT TSB MOD | 3 years | 4-8 | |
| TSB + PHENOL RED | For general use and for the culture of fastidious organisms and not with pH indicator | 1157 B.501157 B.501157 B.501157 B.501157 B.501157 B.501157 B.501157 B.501157 B.501157 B.50 | 50 glass tubes | 210 days | 8-25 | |
| TSB + PHENOL RED | For general use and for the culture of fastidious organisms and not with pH indicator | 1157 B | 10 glass tubes | 210 days | 8-25 | |
| TSB + POLIMIXIN ISO 21871 | For the selective enrichment of B. Cereus | 20998 | 10 glass tubes | 120 days | 4-8 | |
| TSB + POLIMIXIN ISO 21871 | For the selective enrichment of B. Cereus | 20999 | 10 glass tubes 2X | 120 days | 4-8 | |
| TSC ISO 7937 | For the determination and enumeration of Clostridium perfringens | 20509 | 20 plates 90 mm | 180 days | 4-8 | |
| TSC ISO 7937 | For the determination and enumeration of Clostridium perfringens | 20666 | 40 plates 60 mm | 180 days | 4-8 | |
| TSC AGAR BASE ISO 7937 | For the determination and enumeration of Clostridium perfringens | 6646 | Dehydrated 500 gr | 3 years | 8-25 | |
| TSC AGAR BASE ISO 7937 | For the determination and enumeration of Clostridium perfringens | 20171 | 10 glass tubes 20 ml | 180 days | 8-25 | |
| TSC AGAR BASE ISO 7937 | For the determination and enumeration of Clostridium perfringens | 20665 | 10 glass tubes 22 ml | 180 days | 8-25 | |
| TSC AGAR BASE ISO 7937 | For the determination and enumeration of Clostridium perfringens | 20743 | 4 bottles 200 ml | 180 days | 8-25 | |

| | | | | | | |
|--|---|--------|--|----------|------|--|
| TSC AGAR BASE ISO 7937 | For the determination and enumeration of Clostridium perfringens | 20809 | 4 bottles 100 ml | 180 days | 8-25 | |
| TSC AGAR BASE ISO 7937 – REQUIRED SUPPLEMENT | | 6303 | 10 vials x 500 ml / CLOSTRIDIUM PERFRINGENS SUPPLEMENT | 3 years | 4-8 | |
| TSYEA (ISO 11290) | Media for the confirmation of Listeria monocytogenes | 20011 | 20 plates 90 mm | 240 days | 8-25 | |
| TSYEA (ISO 11290) | Media for the confirmation of Listeria monocytogenes | 20012 | 4 bottles 250 ml | 240 days | 8-25 | |
| TSYEA (ISO 11290) | Media for the confirmation of Listeria monocytogenes | 20017 | 4 bottles 100 ml | 240 days | 8-25 | |
| TSYEA (ISO 11290) | Media for the confirmation of Listeria monocytogenes | 20655 | 10 glass tubes 22 ml | 240 days | 8-25 | |
| TSYEA (ISO 11290) | Media for the confirmation of Listeria monocytogenes | 6629 | Dehydrated 500 gr | 3 years | 8-25 | |
| TSYEB (ISO 11290) | Media for the confirmation of Listeria monocytogenes | 20205 | 10 glass tubes | 240 days | 8-25 | |
| TSYEB (ISO 11290) | Media for the confirmation of Listeria monocytogenes | 20455 | 4 bottles 100 ml | 240 days | 8-25 | |
| TSYEB (ISO 11290) | Media for the confirmation of Listeria monocytogenes | 6630 | Dehydrated 500 gr | 3 years | 8-25 | |
| TT HAJNA (USDA-FSIS) | For the selective enrichment of <i>Salmonella</i> spp. in food | 1103 | 10 glass tubes | 90 days | 4-8 | |
| UREA AGAR (ISO 6579) | For the determination of urease activity | 1133 | 10 glass tubes slant | 210 days | 8-25 | |
| UREA AGAR (ISO 6579) | For the determination of urease activity | 1833 | 20 polystyrene tubes slant | 210 days | 8-25 | |
| UREA AGAR (ISO 6579) | For the determination of urease activity | 20682 | 40 plates 60 mm | 210 days | 8-25 | |
| UREA AGAR BASE (ISO 6579) | For the determination of urease activity | 6222 | Dehydrated 500 gr | 3 years | 8-25 | |
| UREA AGAR BASE (ISO 6579) – REQUIRED SUPPLEMENT | | 6530 | 1 x 100 ml / UREA 40% STERILE SOLUTION | 210 days | 4-8 | |
| UREA BROTH BASE | For the determination of urease activity | 6223 | Dehydrated 500 gr | 3 years | 8-25 | |
| UREA BROTH BASE – REQUIRED SUPPLEMENT | | 6530 | 1 X 100 ml / UREA 40% STERILE SOLUTION | 210 days | 4-8 | |
| VIOLET RED BILE GLUCOSE (VRBG) ISO 21528 | For the isolation and the enumeration of Enterobacteria | 1076 B | 20 plates 90 mm | 210 days | 8-25 | |
| VIOLET RED BILE GLUCOSE (VRBG) ISO 21528 | For the isolation and the enumeration of Enterobacteria | 4076 B | 40 contact 55 mm | 210 days | 8-25 | |
| VIOLET RED BILE GLUCOSE (VRBG) ISO 21528 | For the isolation and the enumeration of Enterobacteria | 20424 | 10 glass tubes 15 ml | 210 days | 8-25 | |
| VIOLET RED BILE GLUCOSE (VRBG) ISO 21528 | For the isolation and the enumeration of Enterobacteria | 1276 B | 4 bottles 100 ml | 210 days | 8-25 | |
| VIOLET RED BILE GLUCOSE (VRBG) ISO 21528 | For the isolation and the enumeration of Enterobacteria | 20449 | 4 bottles 200 ml | 210 days | 8-25 | |
| VIOLET RED BILE GLUCOSE (VRBG) ISO 21528 | For the isolation and the enumeration of Enterobacteria | 20448 | 4 bottles 500 ml | 210 days | 8-25 | |
| VIOLET RED BILE GLUCOSE (VRBG) ISO 21528 | For the isolation and the enumeration of Enterobacteria | 6111 | Dehydrated 500 gr | 3 years | 8-25 | |
| VIOLET RED BILE GLUCOSE + LEC + TWEEN 80 + HISTIDINE + SODIUM THIOSULPHATE | For the isolation and the enumeration of Enterobacteria | 21319 | 40 contact 55 mm | 180 days | 4-8 | |
| VIOLET RED BILE LACTOSE AGAR + MUG | Terreno selettivo e differenziale per la determinazione dei Coliformi e per la determinazione diretta di <i>E. Coli</i> | 20383 | 20 plates 90 mm | 90 days | 4-8 | |
| VIOLET RED BILE LACTOSE AGAR + MUG | Terreno selettivo e differenziale per la determinazione dei Coliformi e per la determinazione diretta di <i>E. Coli</i> | 6684 | Dehydrated 500 gr | 2 years | 4-8 | |
| VIOLET RED BILE LACTOSE (VRBL) ISO 4831 | For the isolation, the enumeration and the differentiation of Coliforms | 1076 | 20 plates 90 mm | 210 days | 8-25 | |

| | | | | | | |
|--|---|--------|----------------------|----------|------|--|
| VIOLET RED BILE LACTOSE (VRBL) ISO 4831 | For the isolation, the enumeration and the differentiation of Coliforms | 2276 | 40 plates 60 mm | 210 days | 8-25 | |
| VIOLET RED BILE LACTOSE (VRBL) ISO 4831 | For the isolation, the enumeration and the differentiation of Coliforms | 4076 | 40 contact 55mm | 210 days | 8-25 | |
| VIOLET RED BILE LACTOSE (VRBL) ISO 4831 | For the isolation, the enumeration and the differentiation of Coliforms | 1176 | 10 glass tubes 15 ml | 210 days | 8-25 | |
| VIOLET RED BILE LACTOSE (VRBL) ISO 4831 | For the isolation, the enumeration and the differentiation of Coliforms | 1276 | 4 bottles 100 ml | 210 days | 8-25 | |
| VIOLET RED BILE LACTOSE (VRBL) ISO 4831 | For the isolation, the enumeration and the differentiation of Coliforms | 20819 | 4 bottles 200 ml | 210 days | 8-25 | |
| VIOLET RED BILE LACTOSE (VRBL) ISO 4831 | For the isolation, the enumeration and the differentiation of Coliforms | 6110 | Dehydrated 500 gr | 3 years | 8-25 | |
| VIOLET RED BILE LACTOSE + LEC + TWEEN 80 | For the isolation, the enumeration and the differentiation of Coliforms | 20122 | 40 contact 55 mm | 210 days | 4-8 | |
| VIOLET RED BILE LACTOSE GLUCOSE (VRBLG) (EP) | For the detection of and enumerate the Enterobacteria | 2 | 20 plates 90 mm | 210 days | 8-25 | |
| WL NUTRIENT | For the culture of yeasts and fungi in the beer fermentation processes | 21344 | 20 plates 90 mm | 180 days | 4-8 | |
| WL NUTRIENT | For the culture of yeasts and fungi in the beer fermentation processes | 20417 | 40 plates 60 mm | 180 days | 4-8 | |
| WL NUTRIENT | For the culture of yeasts and fungi in the beer fermentation processes | 6683 | Dehydrated 500 gr | 3 years | 4-8 | |
| WL NUTRIENT + CICLOEXIMIDE | For the isolation of Brettanomyces spp. | 6867 | Dehydrated 500 gr | 3 years | 4-8 | |
| WL NUTRIENT DIFFERENTIAL ISO 10718 | For the isolation of the bacteria in the beer fermentation processes | 21345 | 20 plates 90 mm | 180 days | 4-8 | |
| WL NUTRIENT DIFFERENTIAL ISO 10718 | For the isolation of the bacteria in the beer fermentation processes | 20752 | 40 plates 60 mm | 180 days | 4-8 | |
| WL NUTRIENT DIFFERENTIAL ISO 10718 | For the isolation of the bacteria in the beer fermentation processes | 20855 | 4 bottles 100 ml | 180 days | 4-8 | |
| WL NUTRIENT DIFFERENTIAL ISO 10718 | For the isolation of the bacteria in the beer fermentation processes | 6390 | Dehydrated 500 gr | 3 years | 4-8 | |
| WORT AGAR | For the culture of yeasts and molds | 20439 | 20 plates 90 mm | 180 days | 8-25 | |
| WORT AGAR | For the culture of yeasts and molds | 6225 | Dehydrated 500 gr | 3 years | 8-25 | |
| XLD (EP) | For the isolation of Enterobacteriaceae, especially Salmonella spp. and Shigella spp. | 1027 | 20 plates 90 mm | 180 days | 8-25 | |
| XLD (EP) | For the isolation of Enterobacteriaceae, especially Salmonella spp. and Shigella spp. | 1227 | 4 bottles 100 ml | 180 days | 8-25 | |
| XLD (EP) | For the isolation of Enterobacteriaceae, especially Salmonella spp. and Shigella spp. | 6116 | Dehydrated 500 gr | 3 years | 8-25 | |
| XLD (ISO 6579) | For the isolation of Salmonella spp. | 1027 A | 20 plates 90 mm | 180 days | 8-25 | |
| XLD (ISO 6579) | For the isolation of Salmonella spp. | 1227 A | 4 bottles 100 ml | 180 days | 8-25 | |
| XLD (ISO 6579) | For the isolation of Salmonella spp. | 20277 | 10 plates 150 mm | 180 days | 8-25 | |
| XLD (ISO 6579) | For the isolation of Salmonella spp. | 6219 | Dehydrated 500 gr | 3 years | 8-25 | |
| XLT4 | For the isolation of pathogenic Enterobacteriaceae, especially Salmonella spp. | 1039 A | 20 plates 90 mm | 180 days | 8-25 | |
| XLT4 AGAR BASE | For the isolation of pathogenic Enterobacteriaceae, especially Salmonella spp. | 6635 | Dehydrated 500 gr | 3 years | 8-25 | |

| | | | | | | |
|---|--|-------|--|----------|------|--|
| XYLOSE (D) BROTH ISO 11290 | For the confirmation of Listeria spp. | 20209 | 10 tubes 10 ml | 180 days | 8-25 | |
| YEAST EXTRACT | For use in microbiology | 6418 | Dehydrated 500 gr | 3 years | 8-25 | |
| YEAST EXTRACT AGAR (ISO 6222) | For a broad spectrum growth of bacteria, yeasts and fungi | 20141 | 20 plates 90 mm | 270 days | 8-25 | |
| YEAST EXTRACT AGAR (ISO 6222) | For a broad spectrum growth of bacteria, yeasts and fungi | 20024 | 4 bottles 100 ml | 270 days | 8-25 | |
| YEAST EXTRACT AGAR (ISO 6222) | For a broad spectrum growth of bacteria, yeasts and fungi | 20614 | 4 bottles 250 ml | 270 days | 8-25 | |
| YEAST EXTRACT AGAR (ISO 6222) | For a broad spectrum growth of bacteria, yeasts and fungi | 20675 | 4 bottles 500 ml | 270 days | 8-25 | |
| YEAST EXTRACT AGAR (ISO 6222) | For a broad spectrum growth of bacteria, yeasts and fungi | 6117 | Dehydrated 500 gr | 3 years | 8-25 | |
| YEAST EXTRACT GLUCOSE CHLORAMPHENICOL AGAR (YGC) (ISO 7954) | For the isolation of yeast and fungi in the food | 20380 | 20 plates 90 mm | 180 days | 4-8 | |
| YEAST EXTRACT GLUCOSE CHLORAMPHENICOL AGAR (YGC) (ISO 7954) | For the isolation of yeast and fungi in the food | 20161 | 4 bottles 100 ml | 180 days | 4-8 | |
| YEAST EXTRACT GLUCOSE CHLORAMPHENICOL AGAR (YGC) (ISO 7954) | For the isolation of yeast and fungi in the food | 6291 | Dehydrated 500 gr | 3 years | 8-25 | |
| YEAST NITROGEN BASE WITHOUT AMMONIUM SULFATE | For the classification of the yeasts | 6250 | Dehydrated 500 gr | 3 years | 8-25 | |
| YERSINIA SELECTIVE | For the isolation of Yersinia enterocolitica | 1028 | 20 plates 90 mm | 210 days | 4-8 | |
| YERSINIA AGAR BASE | For the isolation of Yersinia enterocolitica | 6636 | Dehydrated 500 gr | 3 years | 8-25 | |
| YERSINIA AGAR BASE – REQUIRED SUPPLEMENT | | 6399 | 5 + 5 fiale x 500 ml / YERSINIA SELECTIVE SUPPLEMENT | 2 years | 4-8 | |
| YERSINIA ITC BROTH ISO 10273 | For the selective enrichment of Yersinia enterocolitica | 20270 | 10 glass tubes 10 ml | 180 days | 4-8 | |
| YERSINIA ITC BROTH ISO 10273 | For the selective enrichment of Yersinia enterocolitica | 20776 | 10 glass tubes 9 ml | 180 days | 4-8 | |
| YERSINIA ITC BROTH ISO 10273 | For the selective enrichment of Yersinia enterocolitica | 20878 | 4 bottles 90 ml | 180 days | 4-8 | |
| YERSINIA ITC BROTH BASE ISO 10273 | For the selective enrichment of Yersinia enterocolitica | 6715 | Dehydrated 500 gr | 3 years | 8-25 | |
| YERSINIA ITC BROTH BASE ISO 10273 – REQUIRED SUPPLEMENT | | 6394 | 5 + 5 vials x 500 ml / ITK SUPPLEMENT | 2 years | 4-8 | |
| YERSINIA PBS BROTH ISO 10273 | For the selective enrichment of Yersinia enterocolitica | 20879 | 10 glass tubes 9 ml | 180 days | 8-25 | |
| YERSINIA PBS BROTH ISO 10273 | For the selective enrichment of Yersinia enterocolitica | 20775 | 4 bottles 90 ml | 180 days | 8-25 | |
| YERSINIA PBS BROTH ISO 10273 | For the selective enrichment of Yersinia enterocolitica | 6716 | Dehydrated 500 gr | 3 years | 8-25 | |
| YPD BROTH | For the maintenance and the growth of yeasts in molecular biology procedures | 6664 | Dehydrated 500 gr | 3 years | 8-25 | |
| TRIPLE WRAPPED GAMMA IRRADIATIED PLATES | | | | | | |
| PLATE COUNT AGAR (PCA) (APHA) ISO 4833:2003 | For the total aerobic bacterial count in food, water, air and surfaces | 4447 | 40 contact 55 mm | 240 days | 8-25 | |
| PLATE COUNT AGAR (PCA) (APHA) ISO 4833:2003 | For the total aerobic bacterial count in food, water, air and surfaces | 20254 | 20 contact 90 mm | 240 days | 8-25 | |
| PCA + TWEEN 80 + LECITHIN | For the total aerobic bacterial count in food, water, air and surfaces | 4411 | 40 contact 55 mm | 240 days | 4-8 | |
| PCA + TWEEN 80 + LECITHIN | For the total aerobic bacterial count in food, water, air and surfaces | 20255 | 20 contact 90 mm | 240 days | 4-8 | |
| SABOURAUD DEXTROSE | For the culture of yeasts and fungi | 4417 | 40 contact 55 mm | 210 days | 8-25 | |

| | | | | | | |
|--|--|--------|-----------------------|----------|------|--|
| SABOURAUD DEXTROSE | For the culture of yeasts and fungi | 20801 | 20 plates 90 mm | 210 days | 8-25 | |
| SABOURAUD DEXTROSE | For the culture of yeasts and fungi | 4407 | 20 plates 90 mm 25 ml | 210 days | 8-25 | |
| SABOURAUD DEXTROSE | For the culture of yeasts and fungi | 20252 | 20 contact 90 mm | 210 days | 8-25 | |
| SAB + CAF 50 | For the isolation of yeast and fungi | 4418 A | 40 contact 55 mm | 180 days | 4-8 | |
| SAB + LECITHIN + TWEEN 80 | For the culture of yeasts and fungi | 4418 | 40 contact 55 mm | 180 days | 4-8 | |
| SAB + LECITHIN + TWEEN 80 | For the culture of yeasts and fungi | 20253 | 20 contact 90 mm | 180 days | 4-8 | |
| SAB + CAF50 + LECITHIN + TWEEN 80 | For the isolation of yeast and fungi | 4418 B | 40 contact 55 mm | 180 days | 4-8 | |
| TSA (EP) | For the culture of fastidious bacteria and not | 4425 | 40 contact 55 mm | 240 days | 8-25 | |
| TSA (EP) | For the culture of fastidious bacteria and not | 4401 | 20 plates 90 mm 25 ml | 240 days | 8-25 | |
| TSA (EP) | For the culture of fastidious bacteria and not | 20152 | 20 plates 90 mm | 240 days | 8-25 | |
| TSA + LECITHIN + TWEEN 80 | For the culture of fastidious bacteria and not | 4425 A | 40 contact 55 mm | 210 days | 4-8 | |
| TSA + LECITHIN + TWEEN 80 | For the culture of fastidious bacteria and not | 4403 | 20 plates 90 mm 25 ml | 210 days | 4-8 | |
| TSA + LECITHIN + TWEEN 80 + PENASI 20,000 u.i./l | For the culture of fastidious bacteria and not | 20072 | 40 contact 55 mm | 90 days | 4-8 | |
| TSA + LECITHIN + TWEEN 80 + SODIUM THIOSULFATE + HISTIDINE | For the culture of fastidious bacteria and not | 4425 B | 40 contact 55 mm | 210 days | 4-8 | |
| PLATES 90 mm 2 SECTORS | | | | | | |
| CLED / MAC CONKEY | For the presumptive detection and differentiation of organisms causing urinary tract infections - Isolation and differentiation of Enterobacteria | 4203 | 20 plates | 240 days | 4-8 | |
| DERMATOPHYTES / SABOURAUD | Isolation of Dermatophytes - Yeasts and fungi culture | 4294 | 20 plates | 180 days | 4-8 | |
| HEKTOEN / SS | For the isolation and the differentiation of Gram-negative enteric bacteria - For the isolation of Salmonella and Shigella | 4208 | 20 plates | 210 days | 8-25 | |
| SS / MAC CONKEY | For the isolation of Salmonella and Shigella Isolation and differentiation of Enterobacteria | 4220 | 20 plates | 210 days | 8-25 | |
| PLATES 90 mm 3 SECTORS | | | | | | |
| FUNGISCREEN | Yeasts and fungi culture. Isolation of Dermatophytes - Malassezia spp. Culture | 4304 | 20 plates | 180 days | 4-8 | |
| MAC CONKEY / MANNITOL SALT / SABOURAUD | Isolation and differentiation of Enterobacteria - Isolation of S. aureus - Yeasts and fungi culture | 4306 | 20 plates | 180 days | 4-8 | |
| MAC CONKEY / SS / HEKTOEN | Isolation and differentiation of Enterobacteria - For the isolation of Salmonella and Shigella - For the isolation and the differentiation of Gram-negative enteric bacteria | 4305 | 20 plates | 180 days | 4-8 | |
| SABOURAUD / MANNITOL SALT / DESOXYCHOLATE | Yeasts and fungi culture - Isolation of S. aureus - Coliforms isolation and counting | 4301 | 20 plates | 180 days | 4-8 | |
| ANIMAL BLOOD AND DERIVATIVES | | | | | | |
| BLOOD, HORSE DEBRIFINATED STERILE | | 1501 | 25 ml | 30 days | 2-8 | |
| BLOOD, HORSE DEBRIFINATED STERILE | | 1451 | 50 ml | 30 days | 2-8 | |
| BLOOD, HORSE DEBRIFINATED STERILE | | 1452 | 100 ml | 30 days | 2-8 | |

| | | | | | | |
|--|--|------|---------|----------|-----|--|
| BLOOD, HORSE DEBRIFINATED STERILE | | 1453 | 250 ml | 30 days | 2-8 | |
| BLOOD, HORSE DEBRIFINATED STERILE | | 1454 | 500 ml | 30 days | 2-8 | |
| BLOOD, LISATE HORSE DEBRIFINATED STERILE | | 1481 | 50 ml | 365 days | 2-8 | |
| BLOOD, LISATE HORSE DEBRIFINATED STERILE | | 1482 | 100 ml | 365 days | 2-8 | |
| BLOOD, SHEEP DEBRIFINATED STERILE | | 1405 | 25 ml | 30 days | 2-8 | |
| BLOOD, SHEEP DEBRIFINATED STERILE | | 1401 | 50 ml | 30 days | 2-8 | |
| BLOOD, SHEEP DEBRIFINATED STERILE | | 1402 | 100 ml | 30 days | 2-8 | |
| BLOOD, SHEEP DEBRIFINATED STERILE | | 1403 | 250 ml | 30 days | 2-8 | |
| BLOOD, SHEEP DEBRIFINATED STERILE | | 1404 | 500 ml | 30 days | 2-8 | |
| BLOOD, SHEEP DEBRIFINATED STERILE | | 1406 | 1000 ml | 30 days | 2-8 | |
| BLOOD , SHEEP DEFRIFINATED STERILE, IN ALSEVER | | 1421 | 50 ml | 30 days | 2-8 | |
| HORSE SERUM | | 1491 | 50 ml | 365 days | 2-8 | |
| HORSE SERUM | | 1497 | 100 ml | 365 days | 2-8 | |
| HORSE SERUM | | 1496 | 500 ml | 365 days | 2-8 | |